

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/00		A2	(11) International Publication Number: WO 99/33869
			(43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/US98/27416			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 22 December 1998 (22.12.98)			
(30) Priority Data: 08/998,253 24 December 1997 (24.12.97) US 08/998,255 24 December 1997 (24.12.97) US 09/118,627 17 July 1998 (17.07.98) US 09/118,554 17 July 1998 (17.07.98) US			
(71) Applicant: CORIXA CORPORATION [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).			
(72) Inventors: REED, Steven, G.; 2843 - 122nd Place Northeast, Bellevue, WA 98005 (US). XU, Jiangchun; 15805 South-east 43rd Place, Bellevue, WA 98006 (US).			
(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).			


Published

Without international search report and to be republished upon receipt of that report.

(54) Title: COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

(57) Abstract

Compounds and methods for the treatment and diagnosis of breast cancer are provided. The inventive compounds include polypeptides containing at least a portion of a breast tumor protein. Vaccines and pharmaceutical compositions for immunotherapy of breast cancer comprising such polypeptides, or polynucleotide molecules encoding such polypeptides, are also provided, together with polynucleotide molecules for preparing the inventive polypeptides.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RU	Romania		
CU	Cuba	LC	Saint Lucia	SD	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SE	Sudan		
DE	Germany	LK	Sri Lanka	SG	Sweden		
DK	Denmark	LR	Liberia		Singapore		
EE	Estonia						

COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

TECHNICAL FIELD

The present invention relates generally to compositions and methods for the treatment and diagnosis of breast cancer. The invention is more particularly related to polypeptides comprising at least a portion of a protein that is preferentially expressed in breast tumor tissue and to polynucleotide molecules encoding such polypeptides. Such polypeptides may be used in vaccines and pharmaceutical compositions for treatment of breast cancer. Additionally such polypeptides and polynucleotides may be used in the immunodiagnosis of breast cancer.

BACKGROUND OF THE INVENTION

Breast cancer is a significant health problem for women in the United States and throughout the world. Although advances have been made in detection and treatment of the disease, breast cancer remains the second leading cause of cancer-related deaths in women, affecting more than 180,000 women in the United States each year. For women in North America, the life-time odds of getting breast cancer are now one in eight.

No vaccine or other universally successful method for the prevention or treatment of breast cancer is currently available. Management of the disease currently relies on a combination of early diagnosis (through routine breast screening procedures) and aggressive treatment, which may include one or more of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular breast cancer is often selected based on a variety of prognostic parameters, including an analysis of specific tumor markers. See, e.g., Porter-Jordan and Lippman, *Breast Cancer* 8:73-100 (1994). However, the use of established markers often leads to a result that is difficult to interpret, and the high mortality observed in breast cancer patients indicates that improvements are needed in the treatment, diagnosis and prevention of the disease.

Accordingly, there is a need in the art for improved methods for therapy and diagnosis of breast cancer. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

The present invention provides compounds and methods for immunotherapy of breast cancer. In one aspect, isolated polypeptides are provided comprising at least an immunogenic portion of a breast tumor protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein the breast tumor protein comprises an amino acid sequence encoded by a polynucleotide molecule having a partial sequence selected from the group consisting of (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52 and 55-67, 72, 73, and 89-94, (b) complements of said nucleotide sequences and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

In related aspects, isolated polynucleotide molecules encoding the above polypeptides are provided. In specific embodiments, such polynucleotide molecules have partial sequences provided in SEQ ID NOS: 3, 10, 17, 24, 45-52 and 55-67, 72, 73, and 89-94. The present invention further provides expression vectors comprising the above polynucleotide molecules and host cells transformed or transfected with such expression vectors. In preferred embodiments, the host cells are selected from the group consisting of *E. coli*, yeast and mammalian cells.

In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known breast antigen.

The present invention also provides pharmaceutical compositions comprising at least one of the above polypeptides, or a polynucleotide molecule encoding such a polypeptide, and a physiologically acceptable carrier, together with vaccines comprising at least one or more such polypeptide or polynucleotide molecule in combination with a non-specific immune response enhancer. Pharmaceutical compositions and vaccines comprising one or more of the above fusion proteins are also provided.

In related aspects, pharmaceutical compositions for the treatment of breast cancer comprising at least one polypeptides and a physiologically acceptable carrier are provided, wherein the polypeptide comprises an immunogenic portion of a breast tumor protein or a variant thereof, the breast tumor protein being encoded by a polynucleotide molecule having a partial sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88, (b) complements of said nucleotide sequences, and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. The invention also provides vaccines for the treatment of breast cancer comprising such polypeptides in combination with a non-specific immune response enhancer, together with pharmaceutical compositions and vaccines comprising at least one polynucleotide molecule having a partial sequence provided in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88.

In yet another aspect, methods are provided for inhibiting the development of breast cancer in a patient, comprising administering an effective amount of at least one of the above pharmaceutical compositions and/or vaccines.

The present invention also provides methods for immunodiagnosis of breast cancer, together with kits for use in such methods. In one specific aspect of the present invention, methods are provided for detecting breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the inventive polypeptides; and (b) detecting in the sample a protein or polypeptide that binds to the binding agent. In preferred embodiments, the binding agent is an antibody, most preferably a monoclonal antibody.

In related aspects, methods are provided for monitoring the progression of breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent; (c) repeating steps (a) and (b); and comparing the amounts of polypeptide detected in steps (b) and (c).

Within related aspects, the present invention provides antibodies, preferably monoclonal antibodies, that bind to the inventive polypeptides, as well as diagnostic kits

comprising such antibodies, and methods of using such antibodies to inhibit the development of breast cancer.

The present invention further provides methods for detecting breast cancer comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with a first and a second oligonucleotide primer in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In a preferred embodiment, at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID NOS: 1-94.

In a further aspect, the present invention provides a method for detecting breast cancer in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a polynucleotide molecule that encodes one of the above polypeptides; and (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe. Preferably, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID NOS: 1-94.

In related aspects, diagnostic kits comprising the above oligonucleotide probes or primers are provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A and B show the specific lytic activity of a first and a second B511S-specific CTL clone, respectively, measured on autologous LCL transduced with B511s (filled squares) or HLA-A3 (open squares).

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunotherapy and diagnosis of breast cancer. The inventive compositions are generally isolated polypeptides that comprise at least a portion of a breast tumor protein. Also included within the present invention are molecules (such as an antibody or fragment thereof) that bind to the inventive polypeptides. Such molecules are referred to herein as "binding agents."

In particular, the subject invention discloses polypeptides comprising at least a portion of a human breast tumor protein, or a variant thereof, wherein the breast tumor protein includes an amino acid sequence encoded by a polynucleotide molecule including a sequence selected from the group consisting of: nucleotide sequences recited in SEQ ID NOS: 1-94, the complements of said nucleotide sequences, and variants thereof. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising a portion of one of the above breast proteins may consist entirely of the portion, or the portion may be present within a larger polypeptide that contains additional sequences. The additional sequences may be derived from the native protein or may be heterologous, and such sequences may be immunoreactive and/or antigenic.

As used herein, an "immunogenic portion" of a human breast tumor protein is a portion that is capable of eliciting an immune response in a patient afflicted with breast cancer and as such binds to antibodies present within sera from a breast cancer patient. Such immunogenic portions generally comprise at least about 5 amino acid residues, more preferably at least about 10, and most preferably at least about 20 amino acid residues. Immunogenic portions of the proteins described herein may be identified in antibody binding assays. Such assays may generally be performed using any of a variety of means known to those of ordinary skill in the art, as described, for example, in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988. For example, a polypeptide may be immobilized on a solid support (as described below) and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A. Alternatively, a polypeptide may be used to generate

monoclonal and polyclonal antibodies for use in detection of the polypeptide in blood or other fluids of breast cancer patients. Methods for preparing and identifying immunogenic portions of antigens of known sequence are well known in the art and include those summarized in Paul, *Fundamental Immunology*, 3rd ed., Raven Press, 1993, pp. 243-247.

The term "polynucleotide(s)," as used herein, means a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases and includes DNA and corresponding RNA molecules, including HnRNA and mRNA molecules, both sense and anti-sense strands, and comprehends cDNA, genomic DNA and recombinant DNA, as well as wholly or partially synthesized polynucleotides. An HnRNA molecule contains introns and corresponds to a DNA molecule in a generally one-to-one manner. An mRNA molecule corresponds to an HnRNA and DNA molecule from which the introns have been excised. A polynucleotide may consist of an entire gene, or any portion thereof. Operable anti-sense polynucleotides may comprise a fragment of the corresponding polynucleotide, and the definition of "polynucleotide" therefore includes all such operable anti-sense fragments.

The compositions and methods of the present invention also encompass variants of the above polypeptides and polynucleotides. A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. In a preferred embodiment, variant polypeptides differ from an identified sequence by substitution, deletion or addition of five amino acids or fewer. Such variants may generally be identified by modifying one of the above polypeptide sequences, and evaluating the antigenic properties of the modified polypeptide using, for example, the representative procedures described herein. Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity (determined as described below) to the identified polypeptides.

As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids

represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydropathic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

A nucleotide "variant" is a sequence that differs from the recited nucleotide sequence in having one or more nucleotide deletions, substitutions or additions. Such modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983). Nucleotide variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant nucleotide sequences preferably exhibit at least about 70%, more preferably at least about 80% and most preferably at least about 90% identity (determined as described below) to the recited sequence.

The antigens provided by the present invention include variants that are encoded by DNA sequences which are substantially homologous to one or more of the DNA sequences specifically recited herein. "Substantial homology," as used herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the event of cross-species homology, at 45°C with 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS. Such hybridizing DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

Two nucleotide or polypeptide sequences are said to be "identical" if the sequence of nucleotides or amino acid residues in the two sequences is the same when aligned for maximum correspondence as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, more preferably 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins - Matrices for detecting distant relationships: In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) Fast and sensitive multiple sequence alignments on a microcomputer *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) Optimal alignments in linear space *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) The neighbor joining method. A new method for reconstructing phylogenetic trees *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy - the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) Rapid similarity searches of nucleic acid and protein data banks *Proc. Natl. Acad. Sci. USA* 80:726-730.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e. gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by

determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e. the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Also included in the scope of the present invention are alleles of the genes encoding the nucleotide sequences recited herein. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone or in combination with the others, one or more times in a given sequence.

For breast tumor polypeptides with immunoreactive properties, variants may, alternatively, be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. For breast tumor polypeptides useful for the generation of diagnostic binding agents, a variant may be identified by evaluating a modified polypeptide for the ability to generate antibodies that detect the presence or absence of breast cancer. Such modified sequences may be prepared and tested using, for example, the representative procedures described herein.

The breast tumor proteins of the present invention, and polynucleotide molecules encoding such proteins, may be isolated from breast tumor tissue using any of a variety of methods well known in the art. Polynucleotide sequences corresponding to a gene (or a portion thereof) encoding one of the inventive breast tumor proteins may be isolated from a breast tumor cDNA library using a subtraction technique as described in detail below. Examples of such DNA sequences are provided in SEQ ID NOS: 1- 94. Partial polynucleotide sequences thus obtained may be used to design oligonucleotide primers for the amplification of full-length polynucleotide sequences in a polymerase chain reaction (PCR), using techniques well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY,

1989). Once a polynucleotide sequence encoding a polypeptide is obtained, any of the above modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983).

The breast tumor polypeptides disclosed herein may also be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Alternatively, any of the above polypeptides may be produced recombinantly by inserting a polynucleotide sequence that encodes the polypeptide into an expression vector and expressing the protein in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a polynucleotide molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as CHO cells. The polynucleotide sequences expressed in this manner may encode naturally occurring polypeptides, portions of naturally occurring polypeptides, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in an isolated, substantially pure form (i.e., the polypeptides are homogenous as determined by amino acid composition and primary sequence analysis). Preferably, the polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. In certain preferred embodiments,

described in more detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known breast tumor antigen, together with variants of such fusion proteins.

A polynucleotide sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate polynucleotide sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a polynucleotide sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a polynucleotide sequence encoding the second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated polynucleotide sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of polynucleotides are located only 5' to the polynucleotide sequence encoding the first polypeptides. Similarly, stop codons require to end translation and transcription termination signals are only present 3' to the polynucleotide sequence encoding the second polypeptide.

Fusion proteins are also provided that comprise a polypeptide of the present invention together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (see, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91 (1997)).

Polypeptides of the present invention that comprise an immunogenic portion of a breast tumor protein may generally be used for immunotherapy of breast cancer, wherein the polypeptide stimulates the patient's own immune response to breast tumor cells. In further aspects, the present invention provides methods for using one or more of the immunoreactive polypeptides encoded by a polynucleotide molecule having a sequence provided in SEQ ID NOS: 1- 94 (or fusion proteins comprising one or more such polypeptides and/or polynucleotides encoding such polypeptides) for immunotherapy of breast cancer in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease. Accordingly, the above immunoreactive polypeptides (or fusion proteins or polynucleotide molecules encoding such polypeptides) may be used to treat breast cancer or to inhibit the development of breast cancer. The polypeptides may be administered either prior to or following surgical removal of primary tumors and/or treatment by administration of radiotherapy and conventional chemotherapeutic drugs.

In these aspects, the polypeptide or fusion protein is generally present within a pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. The vaccines may comprise one or more of such polypeptides and a non-specific immune response enhancer, wherein the non-

specific immune response enhancer is capable of eliciting or enhancing an immune response to an exogenous antigen. Examples of non-specific-immune response enhancers include adjuvants, biodegradable microspheres (e.g., polylactic galactide) and liposomes (into which the polypeptide is incorporated). Pharmaceutical compositions and vaccines may also contain other epitopes of breast tumor antigens, either incorporated into a combination polypeptide (i.e., a single polypeptide that contains multiple epitopes) or present within a separate polypeptide.

Alternatively, a pharmaceutical composition or vaccine may contain polynucleotides encoding one or more of the above polypeptides, such that the polypeptide is generated *in situ*. In such pharmaceutical compositions and vaccines, the polynucleotide may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems contain the necessary polynucleotide sequences for expression in the patient (such as a suitable promoter). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an epitope of a breast tumor cell antigen on its cell surface. In a preferred embodiment, the polynucleotide molecules may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating polynucleotides into such expression systems are well known to those of ordinary skill in the art. The polynucleotides may also be "naked," as described, for example, in published PCT application WO 90/11092, and Ulmer et al., *Science* 259:1745-1749, 1993, reviewed by Cohen, *Science* 259:1691-1692,

1993. The uptake of naked polynucleotides may be increased by coating the polynucleotides onto biodegradable beads, which are efficiently transported into the cells.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunotherapy of other diseases. In general, the pharmaceutical compositions and vaccines may be administered by injection (e.g., intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (e.g., by aspiration) or orally. Between 1 and 10 doses may be administered over a 3-24 week period. Preferably, 4 doses are administered, at an interval of 3 months, and booster administrations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or polynucleotide molecule that is effective to raise an immune response (cellular and/or humoral) against breast tumor cells in a treated patient. A suitable immune response is at least 10-50% above the basal (i.e., untreated) level. In general, the amount of polypeptide present in a dose (or produced *in situ* by the polynucleotide in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 μ g. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.01 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a lipid, a wax and/or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and/or magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactic glycolide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum

hydroxide or mineral oil, and a nonspecific stimulator of immune response, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis*. Such adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ).

Polypeptides disclosed herein may also be employed in adoptive immunotherapy for the treatment of cancer. Adoptive immunotherapy may be broadly classified into either active or passive immunotherapy. In active immunotherapy, treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (for example, tumor vaccines, bacterial adjuvants, and/or cytokines).

In passive immunotherapy, treatment involves the delivery of biologic reagents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (for example, CD8+ cytotoxic T-lymphocyte, CD4+ T-helper, tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells, lymphokine-activated killer cells), B cells, or antigen presenting cells (such as dendritic cells and macrophages) expressing the disclosed antigens. The polypeptides disclosed herein may also be used to generate antibodies or anti-idiotypic antibodies (as in U.S. Patent No. 4,918,164), for passive immunotherapy.

The predominant method of procuring adequate numbers of T-cells for adoptive immunotherapy is to grow immune T-cells *in vitro*. Culture conditions for expanding single antigen-specific T-cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. These *in vitro* culture conditions typically utilize intermittent stimulation with antigen, often in the presence of cytokines, such as IL-2, and non-dividing feeder cells. As noted above, the immunoreactive polypeptides described herein may be used to rapidly expand antigen-specific T cell cultures in order to generate sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage or B-cells, may be pulsed with immunoreactive polypeptides or transfected with a polynucleotide sequence(s), using standard techniques well known in the art. For example, antigen presenting cells may be transfected with a

polynucleotide sequence, wherein said sequence contains a promoter region appropriate for increasing expression, and can be expressed as part of a recombinant virus or other expression system. For cultured T-cells to be effective in therapy, the cultured T-cells must be able to grow and distribute widely and to survive long term *in vivo*. Studies have demonstrated that cultured T-cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (see, for example, Cheever, M., et al, "Therapy With Cultured T Cells: Principles Revisited," *Immunological Reviews*, 157:177, 1997).

The polypeptides disclosed herein may also be employed to generate and/or isolate tumor-reactive T-cells, which can then be administered to the patient. In one technique, antigen-specific T-cell lines may be generated by *in vivo* immunization with short peptides corresponding to immunogenic portions of the disclosed polypeptides. The resulting antigen specific CD8+ CTL clones may be isolated from the patient, expanded using standard tissue culture techniques, and returned to the patient.

Alternatively, peptides corresponding to immunogenic portions of the polypeptides may be employed to generate tumor reactive T cell subsets by selective *in vitro* stimulation and expansion of autologous T cells to provide antigen-specific T cells which may be subsequently transferred to the patient as described, for example, by Chang et al. (*Crit. Rev. Oncol. Hematol.*, 22(3), 213, 1996). Cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as CellPro Incorporated's (Bothell, WA) CEPRATE™ system (see U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

In another embodiment, T-cell and/or antibody receptors specific for the polypeptides can be cloned, expanded, and transferred into other vectors or effector cells for use in adoptive immunotherapy.

In a further embodiment, syngeneic or autologous dendritic cells may be

pulsed with peptides corresponding to at least an immunogenic portion of a polypeptide disclosed herein. The resulting antigen-specific dendritic cells may either be transferred into a patient, or employed to stimulate T cells to provide antigen-specific T cells which may, in turn, be administered to a patient. The use of peptide-pulsed dendritic cells to generate antigen-specific T cells and the subsequent use of such antigen-specific T cells to eradicate tumors in a murine model has been demonstrated by Cheever et al, *Immunological Reviews*, 157:177, 1997).

Additionally, vectors expressing the disclosed polynucleotides may be introduced into stem cells taken from the patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

Polypeptides of the present invention may also, or alternatively, be used to generate binding agents, such as antibodies or fragments thereof, that are capable of detecting metastatic human breast tumors. Binding agents of the present invention may generally be prepared using methods known to those of ordinary skill in the art, including the representative procedures described herein. Binding agents are capable of differentiating between patients with and without breast cancer, using the representative assays described herein. In other words, antibodies or other binding agents raised against a breast tumor protein, or a suitable portion thereof, will generate a signal indicating the presence of primary or metastatic breast cancer in at least about 20% of patients afflicted with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without primary or metastatic breast cancer. Suitable portions of such breast tumor proteins are portions that are able to generate a binding agent that indicates the presence of primary or metastatic breast cancer in substantially all (*i.e.*, at least about 80%, and preferably at least about 90%) of the patients for which breast cancer would be indicated using the full length protein, and that indicate the absence of breast cancer in substantially all of those samples that would be negative when tested with full length protein. The representative assays described below, such as the two-antibody sandwich assay, may generally be employed for evaluating the ability of a binding agent to detect metastatic human breast tumors.

The ability of a polypeptide prepared as described herein to generate antibodies capable of detecting primary or metastatic human breast tumors may generally be evaluated by raising one or more antibodies against the polypeptide (using, for example, a representative method described herein) and determining the ability of such antibodies to detect such tumors in patients. This determination may be made by assaying biological samples from patients with and without primary or metastatic breast cancer for the presence of a polypeptide that binds to the generated antibodies. Such test assays may be performed, for example, using a representative procedure described below. Polypeptides that generate antibodies capable of detecting at least 20% of primary or metastatic breast tumors by such procedures are considered to be useful in assays for detecting primary or metastatic human breast tumors. Polypeptide specific antibodies may be used alone or in combination to improve sensitivity.

Polypeptides capable of detecting primary or metastatic human breast tumors may be used as markers for diagnosing breast cancer or for monitoring disease progression in patients. In one embodiment, breast cancer in a patient may be diagnosed by evaluating a biological sample obtained from the patient for the level of one or more of the above polypeptides, relative to a predetermined cut-off value. As used herein, suitable "biological samples" include blood, sera and urine.

The level of one or more of the above polypeptides may be evaluated using any binding agent specific for the polypeptide(s). A "binding agent," in the context of this invention, is any agent (such as a compound or a cell) that binds to a polypeptide as described above. As used herein, "binding" refers to a noncovalent association between two separate molecules (each of which may be free (*i.e.*, in solution) or present on the surface of a cell or a solid support), such that a "complex" is formed. Such a complex may be free or immobilized (either covalently or noncovalently) on a support material. The ability to bind may generally be evaluated by determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind" in the context of the present invention when the binding constant for complex formation

exceeds about 10^3 L/mol. The binding constant may be determined using methods well known to those of ordinary skill in the art.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome with or without a peptide component, an RNA molecule or a peptide. In a preferred embodiment, the binding partner is an antibody, or a fragment thereof. Such antibodies may be polyclonal, or monoclonal. In addition, the antibodies may be single chain, chimeric, CDR-grafted or humanized. Antibodies may be prepared by the methods described herein and by other methods well known to those of skill in the art.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding partner to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In a preferred embodiment, the assay involves the use of binding partner immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a second binding partner that contains a reporter group. Suitable second binding partners include antibodies that bind to the binding partner/polypeptide complex. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding partner after incubation of the binding partner with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding partner is indicative of the reactivity of the sample with the immobilized binding partner.

The solid support may be any material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as

adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a second antibody (containing a reporter group) capable of binding to a different site on the polypeptide is added. The amount of second antibody that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior

to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20TM. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of antibody to reporter group may be achieved using standard methods known to those of ordinary skill in the art.

The second antibody is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound second antibody is then removed and bound second antibody is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the

cut-off value is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without breast cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for breast cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for breast cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the antibody is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized antibody as the sample passes through the membrane. A second, labeled antibody then binds to the antibody-polypeptide complex as a solution containing the second antibody flows through the membrane. The detection of bound second antibody may then be performed as described above. In the strip test format, one end of the membrane to which antibody is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second antibody and to the area of immobilized antibody. Concentration of second antibody at the area of immobilized antibody indicates the presence of breast cancer. Typically, the concentration of second antibody at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of antibody immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody

sandwich assay, in the format discussed above. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the antigens or antibodies of the present invention. The above descriptions are intended to be exemplary only.

In another embodiment, the above polypeptides may be used as markers for the progression of breast cancer. In this embodiment, assays as described above for the diagnosis of breast cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, breast cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, breast cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Antibodies for use in the above methods may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.*

6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Monoclonal antibodies of the present invention may also be used as therapeutic reagents, to diminish or eliminate breast tumors. The antibodies may be used on their own (for instance, to inhibit metastases) or coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitzer), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (e.g., U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (e.g., U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (e.g., U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Diagnostic reagents of the present invention may also comprise polynucleotide sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify breast tumor-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a breast tumor protein of the present invention. The presence of the amplified

cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a breast tumor protein of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from SEQ ID NOS: 1-94. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule having a partial sequence provided in SEQ ID NOS: 1-94. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis *et al. Ibid*; Ehrlich, *Ibid*). Primers or probes may thus be used to detect breast tumor-specific sequences in biological samples, including blood, urine and/or breast tumor tissue.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

ISOLATION AND CHARACTERIZATION OF BREAST TUMOR POLYPEPTIDES

This Example describes the isolation of breast tumor polypeptides from a breast tumor cDNA library.

A human breast tumor cDNA expression library was constructed from a pool of breast tumor poly A⁺ RNA from three patients using a Superscript Plasmid System for

cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, breast tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly-A⁺ RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BstX I adaptors (Invitrogen, Carlsbad, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen, Carlsbad, CA) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by electroporation.

Using the same procedure, a normal human breast cDNA expression library was prepared from a pool of four normal breast tissue specimens. The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis. The breast tumor library contained 1.14×10^7 independent colonies, with more than 90% of clones having a visible insert and the average insert size being 936 base pairs. The normal breast cDNA library contained 6×10^6 independent colonies, with 83% of clones having inserts and the average insert size being 1015 base pairs. Sequencing analysis showed both libraries to contain good complex cDNA clones that were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination sequencing.

cDNA library subtraction was performed using the above breast tumor and normal breast cDNA libraries, as described by Hara *et al.* (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a breast tumor-specific subtracted cDNA library was generated as follows. Normal breast cDNA library (70 μ g) was digested with EcoRI, NotI, and SfiI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100 μ l of H₂O, heat-denatured and mixed with 100 μ l (100 μ g) of Photoprobe biotin (Vector Laboratories, Burlingame, CA), the resulting mixture was irradiated with a 270 W sunlamp

on ice for 20 minutes. Additional Photoprobe biotin (50 μ l) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23 μ l H₂O to form the driver DNA.

To form the tracer DNA, 10 μ g breast tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5 μ l H₂O. Tracer DNA was mixed with 15 μ l driver DNA and 20 μ l of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12 μ l H₂O, mixed with 8 μ l driver DNA and 20 μ l of 2 x hybridization buffer, and subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol resistant pBCSK* (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a breast tumor specific subtracted cDNA library.

To analyze the subtracted cDNA library, plasmid DNA was prepared from 100 independent clones, randomly picked from the subtracted breast tumor specific library and characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA). Thirty-eight distinct cDNA clones were found in the subtracted breast tumor-specific cDNA library. The determined 3' cDNA sequences for 14 of these clones are provided in SEQ ID NO: 1-14, with the corresponding 5' cDNA sequences being provided in SEQ ID NO: 15-28, respectively. The determined one strand (5' or 3') cDNA sequences for the remaining clones are provided in SEQ ID NO: 29-52. Comparison of these cDNA sequences with known sequences in the gene bank using the EMBL and GenBank databases (Release 97) revealed no significant homologies to the sequences provided in SEQ ID NO: 3, 10, 17, 24 and 45-52. The sequences provided in SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-41, 43 and 44 were found to show at least some degree of

homology to known human genes. The sequence of SEQ ID NO: 42 was found to show some homology to a known yeast gene.

Data was analyzed using Synteni provided GEMTOOLS Software. Twenty one distinct cDNA clones were found to be over-expressed in breast tumor and expressed at low levels in all normal tissues tested. The determined partial cDNA sequences for these clones are provided in SEQ ID NO: 53- 73. Comparison of the sequences of SEQ ID NO: 53, 54, and 68-71 with those in the gene bank as described above, revealed some homology to previously identified human genes. No significant homologies were found to the sequences of SEQ ID NO: 55-67, 72 (referred to as JJ 9434,7117), and 73 (referred to as B535S).

In a further experiment, cDNA fragments analyzed by DNA microarray were obtained from two subtraction libraries derived by conventional subtraction, as described above. In one instance the tester was derived from primary breast tumors. In the second instance, a metastatic breast tumor was employed as the tester. Drivers consisted of normal breast.

cDNA fragments from these two libraries were submitted as templates for DNA microarray analysis. DNA chips were analyzed by hybridizing with fluorescent probes derived from mRNA from both tumor and normal tissues. Analysis of the data was accomplished by creating three groups from the sets of probes. The composition of these probe groups, referred to as Breast Tumor/mets, Normal non-breast tissues, and Metastatic breast tumors. Two comparisons were performed using the modified Gemtools analysis. The first comparison was to identify templates with elevated expression in breast tumors. The second was to identify templates not recovered in the first comparison that yielded elevated expression in metastatic breast tumors. An arbitrary level of increased expression (mean of tumor expression versus the mean of normal tissue expression) was set at approximately 2.2.

In the first round of comparison to identify overexpression in breast tumors, two novel gene sequences were identified, hereinafter referred to as B534S and B538S (SEQ ID NO: 89 and 90), and six sequences that showed some degree of homology to previously identified genes (SEQ ID NO: 74-79). Additionally, in a second comparison to identify elevated expression in metastatic breast tumors, five novel sequences were identified, hereinafter referred to as B535S (overexpressed in this analysis as well as what was described

above), B542S, B543S, P501S, and B541S (SEQ ID NO: 73, and 91-94), as well as nine gene sequences that showed some homology to known genes (SEQ ID NO: 80-88). Clone B534S and B538S (SEQ ID NO: 89 and 90) were shown to be overexpressed in both breast tumors and metastatic breast tumors.

Example 2

GENERATION OF HUMAN CD8+ CYTOTOXIC T-CELLS THAT RECOGNIZE ANTIGEN PRESENTING CELLS EXPRESSING BREAST TUMOR ANTIGENS

This Example illustrates the generation of T cells that recognize target cells expressing the antigen B511S, also known as 1016-F8 (SEQ ID NO: 56). Human CD8+ T cells were primed *in-vitro* to the B511S gene product using dendritic cells infected with a recombinant vaccinia virus engineered to express B511S as follows (also see Yee et al., Journal of Immunology (1996) 157 (9):4079-86). Dendritic cells (DC) were generated from peripheral blood derived monocytes by differentiation for 5 days in the presence of 50 µg/ml GM-CSF and 30 µg/ml IL-4. DC were harvested, plated in wells of a 24-well plate at a density of 2×10^5 cells/well and infected for 12 hours with B511S expressing vaccinia at a multiplicity of infection of 5. DC were then matured overnight by the addition of 3 µg/ml CD40-Ligand and UV irradiated at 100µW for 10 minutes. CD8+ T cells were isolated using magnetic beads, and priming cultures were initiated in individual wells (typically in 24 wells of a 24-well plate) using 7×10^5 CD8+ T cells and 1×10^6 irradiated CD8-depleted PBMC; IL-7 at 10 ng/ml was added to cultures at day 1. Cultures were re-stimulated every 7-10 days using autologous primary fibroblasts retrovirally transduced with B511S and the costimulatory molecule B7.1. Cultures were supplemented at day 1 with 15 I.U. of IL-2. Following 4 such stimulation cycles, CD8+ cultures were tested for their ability to specifically recognize autologous fibroblasts transduced with B511S using an interferon-γ Elispot assay (see Lalvani et al J. Experimental Medicine (1997) 186:859-965). Briefly, T cells from individual microcultures were added to 96-well Elispot plates that contained autologous fibroblasts transduced to express either B511S or as a negative control antigen EGFP, and incubated overnight at 37° C; wells also contained IL-12 at 10 ng/ml. Cultures were identified that specifically produced interferon-γ only in response to B511S transduced

fibroblasts; such lines were further expanded and also cloned by limiting dilution on autologous B-LCL retrovirally transduced with B511S. Lines and clones were identified that could specifically recognize autologous B-LCL transduced with B511S but not autologous B-LCL transduced with the control antigens EGFP or HLA-A3. An example demonstrating the ability of human CTL cell lines derived from such experiments to specifically recognize and lyse B511S expressing targets is presented in Figure 1.

Example 3

SYNTHESIS OF POLYPEPTIDES

Polypeptides may be synthesized on an Perkin Elmer/Applied Biosystems Division 430A peptide synthesizer using FMOC chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

CLAIMS

1. An isolated polypeptide comprising an immunogenic portion of a breast protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.
2. An isolated polynucleotide molecule comprising a nucleotide sequence encoding the polypeptide of claim 1.
3. An isolated polynucleotide molecule comprising a sequence provided in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94.
4. An expression vector comprising a polynucleotide molecule according to any one of claims 2 and 3.
5. A host cell transformed with the expression vector of claim 4.
6. The host cell of claim 5 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cell lines.
7. A pharmaceutical composition comprising the polypeptide of claim 1 and a physiologically acceptable carrier.
8. A vaccine comprising the polypeptide of claim 1 and a non-specific immune response enhancer.

9. The vaccine of claim 8 wherein the non-specific immune response enhancer is an adjuvant.

10. A vaccine comprising a polynucleotide molecule of any one of claims 2 and 3 and a non-specific immune response enhancer.

11. The vaccine of claim 10 wherein the non-specific immune response enhancer is an adjuvant.

12. A pharmaceutical composition for the treatment of breast cancer comprising a polypeptide and a physiologically acceptable carrier, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

13. A vaccine for the treatment of breast cancer comprising a polypeptide and a non-specific immune response enhancer, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

14. The vaccine of claim 13 wherein the non-specific immune response enhancer is an adjuvant.

15. A vaccine for the treatment of breast cancer comprising a polynucleotide molecule and a non-specific immune response enhancer, the polynucleotide

molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

16. The vaccine of claim 15, wherein the non-specific immune response enhancer is an adjuvant.

17. A pharmaceutical composition according to claims 7 or 12, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

18. A vaccine according to any one of claims 8, 10, 13 or 15, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

19. A fusion protein comprising at least one polypeptide according to claim 1.

20. A pharmaceutical composition comprising a fusion protein according to claim 19 and a physiologically acceptable carrier.

21. A vaccine comprising a fusion protein according to claim 19 and a non-specific immune response enhancer.

22. The vaccine of claim 21 wherein the non-specific immune response enhancer is an adjuvant.

23. A pharmaceutical composition according to claim 20, for use in manufacture of a medicament for inhibiting the development of breast cancer in a patient.

24. A vaccine according to claim 21, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

25. A method for detecting breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent which is capable of binding to a polypeptide, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-94 under moderately stringent conditions; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting breast cancer in the patient.

26. The method of claim 25 wherein the binding agent is a monoclonal antibody.

27. The method of claim 26 wherein the binding agent is a polyclonal antibody.

28. A method for monitoring the progression of breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent that is capable of binding to a polypeptide, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-94 under moderately stringent conditions;

(b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent;

(c) repeating steps (a) and (b); and

(d) comparing the amount of polypeptide detected in steps (b) and (c) to monitor the progression of breast cancer in the patient.

29. A monoclonal antibody that binds to a polypeptide comprising an immunogenic portion of a breast protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

30. A monoclonal antibody according to claim 29, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

31. The monoclonal antibody of claim 30 wherein the monoclonal antibody is conjugated to a therapeutic agent.

32. A method for detecting breast cancer in a patient comprising:

(a) contacting a biological sample from a patient with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotides is specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NO: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions; and

(b) detecting in the sample a polynucleotide sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting breast cancer.

33. The method of claim 32, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from SEQ ID NOS: 1-94.

34. A diagnostic kit comprising:
- (a) one or more monoclonal antibodies of claim 29; and
 - (b) a detection reagent.
35. A diagnostic kit comprising:
- (a) one or more monoclonal antibodies that bind to a polypeptide encoded by a polynucleotide molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88, complements of said sequences and sequences that hybridize to a sequence of SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, or 74-88 under moderately stringent conditions; and
 - (b) a detection reagent.
36. The kit of claims 34 or 35 wherein the monoclonal antibodies are immobilized on a solid support.
37. The kit of claim 36 wherein the solid support comprises nitrocellulose, latex or a plastic material.
38. The kit of claims 34 or 35 wherein the detection reagent comprises a reporter group conjugated to a binding agent.
39. The kit of claim 38 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.
40. The kit of claim 38 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.
41. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein

comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions.

42. A diagnostic kit of claim 41 wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from SEQ ID NOS: 1-94.

43. A method for detecting breast cancer in a patient, comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide probe specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions; and
- (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe, thereby detecting breast cancer in the patient.

44. The method of claim 43 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from the group consisting of SEQ ID NOS: 1-94.

45. A diagnostic kit comprising an oligonucleotide probe specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences, and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions.

46. The diagnostic kit of claim 45, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from the group consisting of SEQ ID NOS: 1-94.

47. Peripheral blood cells from a patient incubated in the presence of at least one polypeptide of claim 1, such that T cells proliferate, for use in the manufacture of a medicament for treating breast cancer in a patient.

48. The blood cells of claim 47 wherein the T cells is repeated one or more times.

49. A composition for the treatment of breast cancer in a patient, comprising T cells proliferated in the presence of a polypeptide of claim 1, in combination with a pharmaceutically acceptable carrier.

50. An antigen presenting cells incubated in the presence of at least one polypeptide of claim 1, for use in the manufacture of a medicament for treating breast cancer in a patient.

51. The cells of claim 50 wherein the antigen presenting cells are selected from the group consisting of dendritic and macrophage cells.

52. A composition for the treatment of breast cancer in a patient, comprising antigen presenting cells incubated in the presence of a polypeptide of claim 1, in combination with a pharmaceutically acceptable carrier.

1/1




Fig. 1A




Fig. 1B

SEQUENCE LISTING

<110> Corixa Corporation

<120> COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

<130> 210121.446PC

<140> PCT

<141> 1998-12-22

<160> 94

<170> PatentIn Ver. 2.0

<210> 1

<211> 402

<212> DNA

<213> Homo sapiens

<400> 1

```

tttttttttt ttttttaggag aactgaatca aacagatttt attcaacttt ttagatgagg 60
aaaacaaatn atacgaaatn ngtcataaga aatgctttct tataccaeta tctcaaacca 120
ctttcaatat ttacaaaaat gctcacgcag caaatatgaa aagctncaac acttcccttt 180
gttaactctg tgcataatnaat gcaacttttaa canacatata aattttctct gtatctctaaa 240
agttnaatta ctaatttttaa tgaatnttnc caagatnttt attcatatac ttttaagtac 300
cctntggcna tacatacanta tttttctttc ttttttttta cnatnggccca acagctttca 360
ngcagncncn aaaaatctta ccggttaatt acacgggggt gt 402

```

<210> 2

<211> 424

<212> DNA

<213> Homo sapiens

<400> 2

```

tttttttttt ttttttaaaag gtacacattt ctttttcatt ctgtttnatg cagcaaaataa 60
tcctgtggca tcttctctgt gatgggcagc ttgctaaaaat tanactcagg ccccttagct 120
ncatttccaa cttagccacac gctttcaacc nngccnaaca aagaaaatca gttingggta 180
aattctttgc tgganacaaa gaactacatt cctttgtaaa tnatgctttg tttgctctgt 240
gcaaacncag attgaaggga anaagganac ttntggggac ggaaacaact ngnagaagca 300
gganccgccc agggncattt cctcaccatg cttatctctg cnetcacttg cngggcacca 360
ttaaacttgg tgcaaaaggc gcaattgggt nanggaaccc cacaccttcc ttaaaaagca 420
gggc 424

```

<210> 3

<211> 421

<212> DNA

<213> Homo sapiens

<400> 3

```

tttttttttt tttttcccaa tttaaaaaag ctttttctat acttcaatta caccanactt 60
aatnatttca tgagtaaatc ngacattatt atttnaaaat ttgcatattt aaaaatttgn 120
tcantttact ccagactgtt tgcanaatga agggaggatc actcaagngc tgatctcna 180
ctntctgcag tctnctgtcc tgtgcccggn ctaattggatc gacactanat ggacagntcn 240

```

```

cagatcttcc gttcttntcc ctcccccaat ttencacenc tccccctctt nccccgaten 300
tttgggggaca tgntaatctt gcnatcctta aacctgtccc gccanggggc ccnanctcag 320
gggtgggttaa tgctcgncng gcttnttgac cncctgccc ctttntntcc naacccccag 420
c 421

```

```

<210> 4
<211> 423
<212> DNA
<213> Homo sapiens

```

```

<400> 4
tttttttatt ttttttctta ttntntntat ttntntnggt tctgtgtgtg aattagnang 60
tgtgtatcgc tangtacnta tgntngcata tttaacctgt tncctttcca tttttaaaa 120
aaaatctcaa natngtantt ggtnnatggg agtaaanaga gactatngat naattttaac 180
atgggacacng tgaatgtag ccgctnatca nttaaaaact tcattttgaa ggccttttnc 240
cctccnaata aaaaatnccng gccctactgg gtttaagcaac attgcatntc taaagaaacc 300
acatgcanac nagttaaac tgtgnaactgg tcangcaaac cnanntggaa nanaagggnn 360
ttcnccecan ggacantcng aattttttta acaattactn atncceccc ngggggggcc 420
tgt 423

```

```

<210> 5
<211> 355
<212> DNA
<213> Homo sapiens

```

```

<400> 5
acgaccacct natttcgtat ctttcaactc ttttcgaccg gacctcttat tcggaagcgt 60
tcacaggaaga caggtctcaa cttagggtac agatcacgtt atcaacgctc tgggactcgt 120
gcaacctggc acttcaagga agtgacccga tnacgtctag accggccaac acagatctag 180
aggtggccaa ctgatcactg taggagctga ctggcaanan tcaacccggc cccaaccnag 240
agtgaccaan acnaccattn aggatcaccc acaggcactc ctgctcctag ggccaaccna 300
ccaaacggct ggccaatggg ggggtttaat atttgggtta aaaattgatt ttaaa 355

```

```

<210> 6
<211> 423
<212> DNA
<213> Homo sapiens

```

```

<400> 6
tttttttttt tttttggaca ggaagtaaaa tttatttgtn antattaana ggggggcagc 60
acattggaag cctcatgan tgcagggccc gccacttgtc cagagggcca cnattggga 120
tgactctaac cccacagccn tctgggatna gccgcttttc agccaccatn tcttcaaat 180
catcagcatt aaacttggtta aanccccact tctttaagat ntgnatcttc tggcgccag 240
naaaacttgaa ctggccctcg cgcaggccct caatcacatg ctccctgttc tgcagctgtg 300
tgcgnaagga cntaatnact tggccnatgt gaacctggc cacantggcc tggggcttc 360
caaaggcacc tcgcaagcct ntttggance tgnccgcccc ngcacaggga caaatcttg 420
ttt 423

```

```

<210> 7
<211> 410
<212> DNA
<213> Homo sapiens

```

```

<400> 7
ttcgactcgt ctaaaacaaa ccgcttgca aagtngaaa aatttatcaa tggaccaaat 60

```

```

aatgctcata tccnacaagt tggtagccgt tnttatnata aaaaaatgta tnatgctcct 120
nanttgttgt acaataatgt tccaatttng gacnttcggc atctaccctg gttcacctgg 180
gtaaatatca ggcagctttt gatggggcta ggaaagctaa cagtactcga acatgggaaa 240
gaggctctgt tgcgcngtgt anattccgct ttgctcngat ttgtggactt 300
catattgttg tacatcgaga tgaatnngaa gaacttgta actactatca ggatcggtgc 360
tttttnaaa agctnatcac catgttggaa cgggcactng gacttgagcg 410

```

<210> 8
 <211> 274
 <212> DNA
 <213> Homo sapiens

```

<400> 8
tttttttttt tttttagggt atacatatatt ttattataa canatatntg tatatacata 60
taatatatgt gtatatatcc acgtgtgtgt ggtggtatca aaacaacaan aantttagtg 120
atctatatct ntngctcaca tatgcatggg agataccagt aaaaaataag tnaatctcca 180
taatatgttt taaaactcan aaaaatcnga gggactnaaa gaaaacgttn acannatga 240
ttgtngataa tcttgaanaa tnaacnaaac atat 274

```

<210> 9
 <211> 322
 <212> DNA
 <213> Homo sapiens

```

<400> 9
tttttttttt ttttgtgctt tattgcacgg gcnanaactt ctagcactat attaaactca 60
ataaggtgta taagtgtgaa aatccttgcc ttctctttaa tcttaatgna naggcactctg 120
gtttttcacc attaantgta ataattggctn tatgtatttt tatnnatggt cttnatggag 180
ttaaaaaggt ttctctctnt cctcngttat ctaanagtgt tnatcaaaaa tgggtataat 240
atttngttca gtacttttnc ctgcacctat agatgatgn ctgttatttt ttcttcttng 300
cctnnanata tgatggatna ca 322

```

<210> 10
 <211> 425
 <212> DNA
 <213> Homo sapiens

```

<400> 10
tttttttttt tttttattct gcagccatta aatgctgaac actagatnct tttttgtgga 60
gggtcacaaaa taagtacaga atathacaca cgcctgtccc ataaaaagca cagctcccag 120
ttcttatatt acaatatctc tggaaatcca ccttcccttc taatttgact aatatcttctg 180
cttctcaggc agcagcgctt tctggcaacc atagaacca acntgnggac taggtcgggtg 240
ggccaaggat caggaaaacag aanaatggaa gnagcccccn tgacnctatt aancntnaaa 300
actatctnaa ctgctagtgt tcaggcttta aatcatgtaa natacgtgtc ctntttgctg 360
caaccgggaag catcctagat ggtacactct ctcagggtgc caggaaaaga tcccaaatng 420
caggn 425

```

<210> 11
 <211> 424
 <212> DNA
 <213> Homo sapiens

```

<400> 11
tttttttant ttttttance nctnntccnn tntgttgnag ggggtaccaa atttctttat 60
ttaaaggaat ggtacaaatc aaaaaactta atttaatttt tnggtacac ttatagaaaa 120

```

```

ggtaagga accccaacat gcatgcactg ccttggtaac caggggnatc ccccnccggt 180
ntggggaaat tagcccaang cttagctttc attatcactn tccccagggt tntgcttttc 240
aaaaaaatct nccgccnagc cnaatccggg cmtcccacac tggcgcaant tgggcacttg 300
gtcccccnat tctttaangg cttncacctn ctcattcggg tnatgtgtct caattaaatc 360
ccacngatgg ggggtcatct tntcnnttag ccagtttgtg nagttccggt attganaaaa 420
ccan 424

```

<210> 12
 <211> 426
 <212> DNA
 <213> Homo sapiens

```

<400> 12
tttttttttt ttttntctaa aagcttttat ctctgctta cattaccat ctgttctgtc 60
atgtgtgtctg ctttttccac tagagccctt aacaacttaa tcatgggtat ttttaagggt 120
ctaataattc cnaaactggg atcataaata agtctcgctt tnatgcttgt tttctctcta 180
tcacactgtg ttngtgtctt ttnnacatgc tttgtaattt ttggctgaaa gctgaaaaat 240
nacatactgt gttntacaac ctgaggtaan cagcctnta gtgtgaggtt ttatatntta 300
ctggctaaga gctnggcncr gtttnantat tgttgtanct ntatatgcca naggtcttna 360
tttcnctng tgctcttgct tnagtacccc attnttttag gggttcccta naaactctat 420
ctnaat 426

```

<210> 13
 <211> 419
 <212> DNA
 <213> Homo sapiens

```

<400> 13
tttttttttt ttttnagat agactctcac tctttcgccc aggctggagt gcagtggcgc 60
aatcaaggct cactgcaacc tctgccttat aaagcatttn ctaaagggtac aagctaaatt 120
tataaataat ctctncacaa ctaagtata acaaaaatta gttctacctc ataaacnct 180
ggctcagccc tcgnaacaca tttccctggt ctcaactgat gaacactcca naaacagaac 240
anantaagc ttttccaggc ccagaaaagc tcgcgagggg atttgctntg tbtgtgacac 300
acttgccacc ctgtggcagc acagctccac acntgctttg ggccgcattt gcaagttctc 360
tgtaancccc ctgnaagacc cggatcagct gggtngaatc tgcangcnct cttttggca 419

```

<210> 14
 <211> 400
 <212> DNA
 <213> Homo sapiens

```

<400> 14
aancattgac caagggtatc cggaggattg tggctgtcac aggtncggag gcccanaagg 60
ccctcaggaa agcaagagc ttgaaaaatg tctctctgtc atggaagccn aagtgaaggc 120
tcactactgt ccaacaagga tntgcanagg gagatcgcta accttgaga ggccttggc 180
actgcagtcn tcccccantg gcagaaggat gaattcgagg agactctcan atcccttang 240
gaaggtcggt gatnacttgg accgagccct nnaagccaat ntccagaaca agtgtggag 300
aagacaaggc anttcatcga cgccaacccc naccgggccc tnttctcctg ganattgana 360
ggcgccccc gcgccagggc cttaataanc cntgaagctn 400

```

<210> 15
 <211> 395
 <212> DNA
 <213> Homo sapiens

<400> 15
 tgctttgctg cgtccaggaa gattagatng aanaatacat attgatttgc caaatgaaca 60
 agcgagatta gacntactgc anacccatgc aggtcccatc acaagcatg gtgaaataga 120
 tgatgaagca attgtgaagc tatcgatggc ctttnatgga gcagatctga gaaatgtttg 180
 tactgaagca ggtatgttcg caattcgtgc tgatcatgat tttgtagtac aggaagactt 240
 catgaaagcn gtcagaanag tggctnattc tnaaagctgg agtctaatt ggaacnacac 300
 ctntgtattt actgttggan ttttgatgct gcatgacaga ttttgcttan tgtaaaaatn 360
 aagtcaaga aaattatgtt agttttggcc attat 395

<210> 16

<211> 404

<212> DNA

<213> Homo sapiens

<400> 16
 ccaccactaa aatcctggct gagccctacn agtacctgtg cccctccccc aggacgagat 60
 nagggcacac cctttaagtn aggtgacagg tcacctttaa gtgaggacag tcagctnaat 120
 ttacactctt gggcttgagt acctggttct cgtgcccctga ggcgacnctn agccctgcag 180
 ctncatgt a cgtgctgcca atngtcttga tcttctccac gccnctnaac ttgggcttca 240
 gtaggagctg caggcnagaa ngaagcgggt aacagcgcca ctccatagcc gcagcnggc 300
 tgcccctgct tctcaaggag ggggtgtggg ttccctccacc atcgccgcc ttcgaaacac 360
 ntctcanggc ttcctnccg gctnancgca ngacttaagc atgg 404

<210> 17

<211> 360

<212> DNA

<213> Homo sapiens

<400> 17
 ggccagaagc ttctcacaaa ccagtgaagg tggcgagcaa gaaagcctct tagacnagga 60
 gctggcgaga gctgctatct ngatngacng cagaaaccaa ccactaattc agcaaacaca 120
 acctcatacc tnaaccgttct cctttnaatg gccttcggtg tggcgagaca tgggcacgtg 180
 cgtgggagaa catactatt cccctnttcc cggcctacca cctctnctcc cccctctctt 240
 ctctncaatt actnctctcn ctgcttntt ctanacacta ctgctngnt cnaagccng 300
 cccgcaatta cctggcaaaa ctgcgagacc ttcgggcagc gctaaaanaa gcacatttac 360

<210> 18

<211> 316

<212> DNA

<213> Homo sapiens

<400> 18
 atacatatat acatatatga ttttagatag agccatatac ctngaagtag tanatttgtt 60
 tgtgtgtata tgtatgtgtc tactcatctt aaataaacct gtgatagaga tgtaattntg 120
 agccagtttt tcatctgctt aaatnactca ccaagtaact aattaagtn tctttactct 180
 taatgttnag tagtgagatt ctgttgaagg tgatattaaa aaccattcta tattaattaa 240
 cattcatgtt gttttttaaa agcttatttg aaatcnaatt atgattattt ttcataccag 300
 tcgatnttat gtangt 316

<210> 19

<211> 350

<212> DNA

<213> Homo sapiens

<400> 19


```

aagggatgca nataatgctg tgtatgagct tgatggaaaa gaactctgta gtgaaaggg 60
tactattgaa catgctnggg ctcggtcacg aggtggaaga ggtagaggac gatactctga 120
ccgttttagt agtcgcagac ctcgaaatga tagacgaaat gctccacctg taagaacaga 180
anacgtctct atagtggaga atttatcctc aagagtcagc tggcaggttt gtganatac 240
agttttgtag tnttttgatg tggcttttta aaaaagttat gggttactna ttgtattatg 300
ttttataaa agtagttttn aattaatgga tntgatggaa ttgtgtttt 350

```

<210> 20

<211> 367

<212> DNA

<213> Homo sapiens

<400> 20

```

gntnnnncca agatcetnet ntccccnngg gcngccccc cncngtnat naccgggttn 60
ntaanatcnn gccgcncccg aagctctcnc nntgccgaga tgncccttat ncnncnatgn 120
ncaattntga cctnngggcga anaatggcng nngtgtatca gtntcncctc tgnngnctct 180
tagnatctga ccactangac cncctatcct ctcaaacctc gtannngccc ctaattttgt 240
ccaattagtg catgntanag cntcctggcc cagatggcnc ccatatccctg gtncgggttc 300
cgccctacc angncatccn catctactag agcttatccg ctncmtgngg cgcaccggnt 360
cccnct 367

```

<210> 21

<211> 366

<212> DNA

<213> Homo sapiens

<400> 21

```

cccaacacaa tggctctaagt anaactgtat tgcctctgtag tatagttcca cattggcaac 60
ctacaatggg aaaatccata cataagtcag ttacttcctn atgagcttcc tccctctgaa 120
tcctttatct tctgaagaaa gtacacaccc tggtnatgat atctttgaat tgcctctctt 180
tccaggcatc agttggatga ttcatcatgg taattatggc attatcatac tcttcatact 240
tgtcatcaga aaacaccagt tctgcccna gatgagcttg ttctgcagct cttagcaact 300
tgggaatatt cactctagac cagaaaacag tcccgtgtct ccttcatttt ctgaggctta 360
aatttn 366

```

<210> 22

<211> 315

<212> DNA

<213> Homo sapiens

<400> 22

```

acttaatgca atctctggag gataatttgg atcaagaaat aaagaanaaa tgaattagga 60
gaagaatna ctgggtntata ttccaatatt ttagaacttt aanaatgttg actatgatt 120
caatatatt gtnaaaactg agatacangt ttgacctata tctgcatttt gataaataaa 180
cnaatnnatt ctattttaat gtgtgttcag agtcacagca cagactgaaa ctttttttga 240
atacctnaat atcacacttn tnccttnaat gatgtggaag acaatgatga catgccttna 300
gcataataat tcgac 315

```

<210> 23

<211> 202

<212> DNA

<213> Homo sapiens

<400> 23

```

actaatccag tgggtgtnaa ttccattgtg ttgggcaact caggatatta aatttatnat 60

```

ttaaaaatc ccaagagaaa naaactccag gccctgattg tttcactggg gaattttacc 120
 aaatgttnca nnaaganatg acgctgattc tgnnaaatct ttttcagaa atagaggaga 180
 acaccacccg ntctatttta tg 202

<210> 24

<211> 365

<212> DNA

<213> Homo sapiens

<400> 24

ggattttcttg cctttttctc cctttttaag tatcaatgta tgaatccac ctgtaccacc 60
 cttttcgcca tacaaccgct accacatctg gctcctagaa cctgttttgc ttcatagat 120
 ggatctcgsa accnagtgtt nacttctatt ttaaaccaca ttttagcaga tngtttgc 180
 tggttctgtct gtattccacca tggggcctgt acacaccag tgtgggtata gtcaaacaca 240
 gtgacctcca ttgtggccac atggggagacc catnaccna tactgcatcc tgggctgatn 300
 acggcactgc atctnaccg acntgggatt gaacccgggg tgggcagcng aattgaacag 360
 gatca 365

<210> 25

<211> 359

<212> DNA

<213> Homo sapiens

<400> 25

gtttctctgct tcaacagtgc ttggacggaa ccggcgctc gttccccacc ccggccggcc 60
 gcccatagcc agccctcgt cactcttca ccgcaccctc ggactgccc aaggcccccc 120
 cgccnctcc ngcgcnccg agccaccgcc cctcctctn gtcccgccnt 180
 nacaacgct ccactcgca ngttcgcnng aactaccacc nggactcata ngccgcccc 240
 aaccgcoga tcaacctgga gctctncccc ccgaentaa cctttccntg tcttaetac 300
 ntaaccgcc gnttattttg cttnaaaaga acttttccc aatactttct ttcaccnnt 359

<210> 26

<211> 400

<212> DNA

<213> Homo sapiens

<400> 26

agtgaacacg tatatgtgaa aaggagtttg tgannagcta cataaaaaa ttatgatct 60
 ttataatttc caataggata ctcatcagtt ttgaataana gacatattct agagaaacca 120
 ggtttctcgt ttacgatttg aactctcaag agcttgaag ttatcactcc catctcag 180
 acnacnaana aatctnaacn aacngaanac caatgacttt tcttagatct gtcaagaac 240
 ttcagccacg aggaaaaacta tcnccctnaa tactggggac tggaaagaga ggtacagag 300
 aactcacagt aatcatagcc caagatcagc ttgcccgag ctnaagactng tacgatnatt 360
 acctacaggg accacttcac agtnngtnga tnaantgcn 400

<210> 27

<211> 366

<212> DNA

<213> Homo sapiens

<400> 27

gaattttcta gaaactgaag ttactctgt tccaagatat atcttctactg tcttaataca 60
 agggcgctng aatcatagca aatatttca tctttcaact aactttaagt agttntcctg 120
 gaattttaca ttttcagaa aacactcctt tctgtatctg tgaagaaaag tgtgcctcag 180
 gctgtagact gggctgcact ggacacctgc gggggactct ggctnagctn ggacatggct 240

```

agtattgatt ttcctcanac tcagcctgtg tagctntgaa agcatggaac agattacact 300
gcagttnacg tcatcccaca catcttggac tcenagaccg ggggagggtca catagtccgt 360
tatgna 366

```

<210> 28

<211> 402

<212> DNA

<213> Homo sapiens

<400> 28

```

agtgggagcc tcttccttec ccactcagtt ctttacatcc ccgaggcgca gctgggcnaa 60
ggaagtggcc agctcgagcg cctcctgcag gcagccaacg tcttgcctg tggcctgtgc 120
agacacatcc ttgccaccac cttacccgct catcangcct gacacctgct gcacccactc 180
gctngctttt aagccccgat nggctgcatt ctgggggact tgacacagcg ncgtgatctt 240
gccagcctca ttgtccaccg tgaagagcat ggcacaaaagt ctgaggggag tgcattcttga 300
anagcttcaa ggcttcattc agggccttng ctnaggcgcc nctctccatc tccnggaata 360
acnagaggct ggtnggggtg actntcaata aactgcttgc tc 402

```

<210> 29

<211> 175

<212> DNA

<213> Homo sapiens

<400> 29

```

cggacgggca tgaccggtcc ggtcagctgg gtggccagtt tcagttcttc agcagaactg 60
tctcccttct tggggggcga gggcttcctg gggagagga tgagtttggg gcggtactcc 120
ttcagccgct gcacgttggg ctgcaggagc tccgtggact tgttccgcct cctcg 175

```

<210> 30

<211> 360

<212> DNA

<213> Homo sapiens

<400> 30

```

ttgtatttct tatgatctct gatgggttct tctcgaatat gccaaagtga agactttgtg 60
gcagtctcca gatttaaact cagctgagcg tccctttgtt ttcagttcca tgaataactc 120
tggagaggaa cttcacggag aggaagactg ctggagaaga gaagcgtgtt agccccattg 180
aggtctgggg aatcatgtaa agggtaacca gacctcactt ttatgtattt acatcaatga 240
gtctcttcag ggaaccaaac ccagaattcg gtgcacaaag caaacatctt ggtgggattt 300
gataaatgcc ttgggacctg gagtgcctgg cttgtgcaca ggaagagcac cagccgctga 360

```

<210> 31

<211> 380

<212> DNA

<213> Homo sapiens

<400> 31

```

acgctctaag cctgtccacg agctcaatag ggaagcctgt gatgactaca gactttcgca 60
acgctacgcc atggtttatg gatacaatgc tgccctataa cgctacttca ggaagcgccg 120
agggaccnaa tgagactgag ggaagaaaaa aaatctcttt tttctggag gctggcacct 180
gattttgtat cccctctgtnn cagcattncn gaaatacata ggcttatata caatgctctt 240
ttcctgtata ttctctgttc tggctgcacc ccttnttccc gcccccagat tgataagtaa 300
tgaaagtga ctgcagtnag ggtcaangga gactcancat atgtgatgt tccntnataa 360
acttctggtg tgatactttc

```

<210> 32
 <211> 440
 <212> DNA
 <213> Homo sapiens

<400> 32
 gtgtatggga gcccctgact cctcacgtgc ctgactctgt cccttggtcc caggctcaggc 60
 ccacccctgt caccctccacc tgccccagcc cctgcctctg ccccaagtgg gccagctgc 120
 cctcacttctl ggggtggatg atgtgacctt cctnggggga ctgcggaagg gacaaagggt 180
 ccctgaagtc ttacggtcca acatcaggac caagtcctcat ggacatgctg acagggtccc 240
 caggggagac cgtntcanta gggatgtgtg cctggctgtg tacgtgggtg tgcagtgcac 300
 gtganaagca cgtggcggct tctgggggccc atgtttgggg aagggaagtg gccnccacc 360
 cttggagaac ctacgtcccn gtacgccctt gccctggcac agcngcatnc acttcaaggg 420
 caccctttgg ggggtggggt 440

<210> 33
 <211> 345
 <212> DNA
 <213> Homo sapiens

<400> 33
 tattttaaca atgtttatta ttcattttat cctctataga accaccaccc acaccgagga 60
 gattatttgg agtgggtccc aacctagggc ctggactctg aaatctaact ccccacttcc 120
 ctcatcttctg gacttaggtg ggggcattgt tcagtcagaa ctgggtgtctc ctatggatgc 180
 gtgcagaagg aggaactagg cacacacata tgggtggccac acccaggagg gttgattggc 240
 aggtctggaag acaaaagtct cccaataaag gcacttttacc ctcaaagang ggggtgggagt 300
 tggctctgtg ggaatgtgtg tgttgggggt ggggaagant atttc 345

<210> 34
 <211> 440
 <212> DNA
 <213> Homo sapiens

<400> 34
 tgtaattttt ttattggaaa acaaatatac aacttggaaat ggattttgag gcaatttgtg 60
 ccataagcag atttttaagt gctaaacaaa gtttaaaaaag caagtaacaa taaaagaaaa 120
 tgtttctgtt acaggaccag cagtcacaaa aaatagtgtg cgagtaacct gataatacac 180
 ccgttttgcg atagtgcacac ttttaagtac atattgttga ctgtccatag tccacggaga 240
 gttacaactc cacacttcaa caacaacatg ctgacagttc cttaagaaaa ctactttaaa 300
 aaaggcctaa cccagatgtt cctctatttg accaactcca tctnagttaa gatgtgcaga 360
 agggcttana ttttccaga gtaaggccna tgcaacatgt tacttgatca attttctaaa 420
 aaagggtttt aggacaatga 440

<210> 35
 <211> 540
 <212> DNA
 <213> Homo sapiens

<400> 35
 atagatggaa tttattaagc ttttcacatg ttagtagcaca tagttttta tgcattccaaa 60
 gtactaacaa aaactctagc aatcaagaat ggcagcatgt tattttataa caatcaacac 120
 ctgtggcttt taaaatttgg ttttcataag ataattata ctgaagttaa tctagccatg 180
 cttttaaaaa atgcttttagg tcaactccaag cttggcagtt aacatttggc ataaacaata 240
 ataaacaact caaattttaa taataacaa atacaacatt gtaggccata acatataca 300
 gtataaggga aaagggtgta gtgttganta agcagttatt agaatagaat accttggcct 360

ctatgcaaat atgtctagac actttgattc actcagccct gacattcagt tttcaaggt 420
 aggaacacagg tctacagta tcattttaca gtctccaaac cattgaaaac aagtagaaaa 480
 tgatganttgt atttttatta atgcattaca tcttcaagan ttatcaccaa cccctcaggt 540

<210> 36

<211> 555

<212> DNA

<213> Homo sapiens

<400> 36

cttcgtgtgc ttgaaaattg gagcctgccc ctcgcccat aagcccttgt tgggaactga 60
 gaagtgata tggggcccaa nctactggtg ccagaaacaca gagacagcag cccantgcaa 120
 tgctgtcgag cattgcacac gccatgtgtg gaactaggag gaggaatatt ccatcttggc 180
 agaaacacaca gcattgggtt ttttctactt gtgtgtctgg gggaaatgaac gcacagatct 240
 gtttgacctt gttataaaaa tagggctccc ccacctcccc cntttctgtg tnccttattg 300
 tagcantgct gtctgaagg gagccctcan cccctggcag acanancctg ttcagtgtccc 360
 ctttctctc tgctaaatgg atgttgatgc actggaggtc ttttancctg cccctgcatg 420
 gncctctgtg gaggaagana aaactctgct ggcgatgacc acagtttctt gactggangc 480
 cntcaaccct cttggttgaa gccttggtct gaccctgaca tntgcttggg cncctgggtng 540
 gntggggctt ctnaa 555

<210> 37

<211> 280

<212> DNA

<213> Homo sapiens

<400> 37

ccaccgacta taagaactat gccctcgtgt attcctgtac ctgcatcatc caacttttcc 60
 acgtggattt tgcttggatc ttggcaagaa accctaactc cctccagaa acagtggact 120
 ctctaaaaaa tatcctgact tctaataaca ttgatntcaa gaaaatgagc gtacagagacc 180
 aggtgaactg ccccnagctc tcgtaacdag gtctacagg gaggtctgac ccactccatg 240
 ttncctctgc ttcgctttcc cctacccacc ccccgccat 280

<210> 38

<211> 303

<212> DNA

<213> Homo sapiens

<400> 38

catcgagctg gttgtcttct tgccctgccct gtgtcgtaaa atgggggtcc cttactgcat 60
 tatcaaggga aaggcaagac tgggagctct agtccacagg aagacctgca ccactgtcgc 120
 cttcacacag gtgaactcgg aagacaaagg cgcttttgct nagctgggtg aagctatcag 180
 gaccaattac aatgacngat acgatnagat ccgcctncac tggggtagca atgtcctggg 240
 tctaaagtct gtggctcgta tcgccnagct cgaanaggcn aangctaag aacttgccac 300
 taa 303

<210> 39

<211> 300

<212> DNA

<213> Homo sapiens

<400> 39

gactcagcgg ctggtgtctt tcctgtgcac aagcccagca ctccaggctcc caaggcattt 60
 atcaaatccc accaagatnt ttggcttttg caccgaattc tgggttttgt tccttnaag 120
 aactcattga tgaatatnac tnaaagttag gtctgggtac cctttacatg attccccaga 180

cctcanatgg gctaacacgc ttctcttctc cagcagctctt cctntccgtg aagttacctt 240
ccagattgtt acatggaact gaanacaaag ggagcctcag ctngatttaa atctggagca 300

<210> 40

<211> 318

<212> DNA

<213> Homo sapiens

<400> 40

cccaacacaa tggctgagga caaatcagtt ctctgtgacc agacatgaga aggttgccaa 60
tgggctgttg ggcgaaccaag gccttccggg agtcttcgtc ctctatgagc tctcgcccat 120
gatgggtgaag ctgacggaga agcacagggtc ctccaccacac ttcttgacagc gtggtgtgcgc 180
catcatgtggg ggcattgttca cagtggctgg actcatcgat tcgtccatct accactcagc 240
acagccatc cagaaaaaaa ttgatctnng gaagacnacg tagtcacctt cggtncttcc 300
tctgtctctt cttttctcc 318

<210> 41

<211> 302

<212> DNA

<213> Homo sapiens

<400> 41

acttagatgg ggtccgttca ggggatacca gcgttcacat ttttctttt aagaaagggt 60
cttggcctga atgttcccca tccggacaca ggctgcattgt ctctgtnagt gtcaaaagctg 120
ccatnaccat ctggttaacc tactcttact ccacaatgtc tatnttcaat cgagggtctct 180
ataatnagtc cataatgtaa atgcctggcc caagaacntat ggctgtgagt tatccnaggc 240
ccaaacnatt accagacatt cctcttanat tgaaaaacga tntctttccc ttggcaaaaa 300
tc 302

<210> 42

<211> 299

<212> DNA

<213> Homo sapiens

<400> 42

cttaataagt ttaaggccaa ggcccgttcc attcttctag caactgacgt tggcagccga 60
ggtttgagca taacctcatgt aaatgtgggt gtcaactttg acattcttac ccattccaag 120
gattacatcc atcgagtagg tcgaacagct agagctgggc gtcccggaag ggctattact 180
tttgtcacac agtatgatgt ggaactcttc cagcgcatag aacacttnat tgggaagaaa 240
ctaccaggtt ttccaacaca ggatgatgag gttatgatgc tnacggaacg cgtcgctna 299

<210> 43

<211> 305

<212> DNA

<213> Homo sapiens

<400> 43

ccaacaatgt caagacagcc gtctgtgaca tccacctcgt tggcctdaan atggcagtc 60
ccttcatctg caatagcaca gccntccggg agctcttcaa gcgcatctcg gagcagttca 120
ctgcatctgt ccgccggaag gccttctctc actgggtacac aggcgagggc atggacaaga 180
tggagttcac cgaggctgag agcaacatga acgacctcgt ctctnagtat cagcagttacc 240
gggatgccac ccagaaaana ggaggaggat ttgggtnagg aggcgaaga aggaggcctg 300
aggca 305

<210> 44

<211> 399

<212> DNA

<213> Homo sapiens

<400> 44

```

ttctctgtggg ggaaacctga tctcgacnaa attagagaat ttgtcagcg gtatttcggc 60
tggaacacgaa cgaaaaacnga tnaatctctg ttctctgtat taaagcaact cgaatncccg 120
cagacacacg tccnaattga ttctctcttt ngattagcac aacaggggaga aagaanaatgc 180
ttaacgtatt aagagccnga gactaaacag agctttgaca tgtatgcta ggaaagagaa 240
agaagcagcn gcccgcnnaa ttngaagcng ttctctgtgc cntgganaaa gaatttgagc 300
ttctttatta ggccaacgaa aaaccccgaa ananaggcnt tacnatacct tngaaaantc 360
tccngccnna aaaagaaaga agctttcnga ttcttaacc 399

```

<210> 45

<211> 440

<212> DNA

<213> Homo sapiens

<400> 45

```

gcgggagcag aagctaaagc caaagcccaa gagagtggca gtgccagcac tgggtgccagt 60
accagatacca ataacagtgc cagtgcagcag gccagcacca gtgggtggctt cagtgcctggc 120
gccagcctga ccgccactct cacatttggg ctctctcgctg gccttgggtgg agctgggtgcc 180
agcacacagtg gcagctctgg tgctctgggt ttctctctaca agtgagattt taggtatctg 240
ccttgggttct agtggggaca tctggggctt anggggcnng gataaaggagc tggatgattc 300
taggaaggcc cangttggag aangatgtgn anagtgtgcc aagacactgc ttttggcatt 360
ttattctctt ctgtttgtcg gangtcaatt gacctttnna ntctctcta ctgtgtttt 420
canatatngt taatcctgcc 440

```

<210> 46

<211> 472

<212> DNA

<213> Homo sapiens

<400> 46

```

gtctctgaat ttacatttt aaaccttccc ttgacctcac attctcttc gccaccctct 60
gtttctctgt tctctctcac agcaaaaaact gttcaaaaaga gttgttgatt actttcattt 120
ccaactttct accccacttc tccctcaat taactctctt tcatccccat gatgccatta 180
gttggctntt attanagtca ccaaccttat tctccaaaac anaagcaaca aggactttga 240
cttctcagca gcaactcagct ctggtncttg aaacaccccc gtacttgcg attctctcta 300
ctctcaatac atctcttccc cagcctctac tgcctgcttc tctgagttct tcccagggtc 360
ctaggctcag atgtagtgtg gctcaacctt gctacacaaa gnaactctct gaagcctgt 420
aaaaatgtcc atnctgtgcc tgtgagtgat ctncangna naataacaaa tt 472

```

<210> 47

<211> 550

<212> DNA

<213> Homo sapiens

<400> 47

```

ccttctcccg cctggccatc cccagcatgc tcatgtctgt catggagtgg tgggacctatg 60
aggtcgggag ctctctcagt ggtctgtatg aggatggatg acggggactg gtgggaacct 120
ggggggccctg tctgggtgca aggcgacagc tgcctttctt caccaggcat cctcgccatg 180
gtggagctcg gcctcagtc catcgtgtat gaactggcca tcatgtgta catggctccc 240
gcaggcttca gtctggctgc cagtgctcgg gtangaacag ctctgggtgc tggagacatg 300
gaagcaggca cggaagtctc ctaccgttct cctgctgatt acagtgtctt ttgctgtanc 360

```

```

cttcagtgtc ctgctgttaa gctgtaagga tcacntgggg tacattttta ctaccgaccg 420
agaacatcat taactctgtg gctcaggtgg ttccaattta tgctgtttcc cactcttttg 480
aagctcttgc tgctcaggtta cagccaatt ttgaaaagta aacaacytgc ctcggagtgg 540
gaattctgtc                                     550

```

<210> 48

<211> 214

<212> DNA

<213> Homo sapiens

<400> 48

```

agaaggacat aaacaagctg aacctgccca agacgtgtga taccagcttc tcagatccag 60
acaacctcct caacttcaag ctggctatct gtcctgatna gggcttctac nagagtggga 120
agtttgtgtt cagttttaag gtgggccagg gttaccgcga tgatcccccc aaggtgaagt 180
gtgagacnat ggtctatcac ccnaccattg acct                                     214

```

<210> 49

<211> 267

<212> DNA

<213> Homo sapiens

<400> 49

```

atctgcctaa aatttattca aataatgaaa atnaatctgt tttagaatat tcagtccttt 60
agtttttagg acaactatgc acaaatgtac gatggagaat tctttttgga tnaactctag 120
gtngaggaac ttaactcaac cggagctntt gtgaaggtca gaanacagga gagggaaatct 180
tggcaaggaa tggagaacga gtttgcaaat tgcagctaga gtnaatngtt ntaaatggga 240
ctgctnttgt gtctcccgang gaaagtt                                     267

```

<210> 50

<211> 300

<212> DNA

<213> Homo sapiens

<400> 50

```

gactgggtca aagctgcatg aaaccaggcc ctggcagcaa cctgggaatg gctggaggtg 60
ggagagaacc tgactctctt ttccctctcc ctctcccaac attactggaa ctctgtcctg 120
gtggatctct ctgagcttgt ttccctgctg ggtgggacag aggacaaagg agaaggaggag 180
gtctagaaga ggcagccctt ctttgtctct tggggtnaat gagcttgacc tanagttagt 240
ggagagacca anagcctctg atttttaatt tccataanaat gttcnaagta tatntntacc 300

```

<210> 51

<211> 300

<212> DNA

<213> Homo sapiens

<400> 51

```

gggtaaaaac ctgcagcacc cactctggaa aatactgtct ttaattttcc tgaaggtggc 60
ccctattttc tagttggtcc aggattaggg atgtggggta tagggcattt aaactctctc 120
aagcgtctct caagcaccoc cggcctgggg gtnagtttct catcccgcta ctgctgctgg 180
gatcaggttn aataaatgga actcttctct tctggcctcc aaagcagcct aaaaactgag 240
gggctctgtt agagggggacc tccaccctnn ggaagtcoga ggggctnngg aaggggttct 300

```

<210> 52

<211> 267

<212> DNA

<213> Homo sapiens

<400> 52

```

aaaatcaact tcntgcatta atanacanat tctanancag gaagtgaana taattttctg 60
cacctatcaa ggaacnnact tgattgcctc tattnaacan atatatcgag ttnctatact 120
tacctgaata ccnccgcata actctcaacc nanatnctc nccatgacac tcnttcttna 180
atgctantcc cgaattcttc attatatcng tgatgttcgn cctgntnata tatcagcaag 240
gtatgtncen taactgcoga nncaang                                     267

```

<210> 53

<211> 401

<212> DNA

<213> Homo sapiens

<400> 53

```

agsetttagc atcatgtaga agcaaaactgc acctatggct gagatagggt caatgacctt 60
caagattttg tgttttctag ctgtccagga aaagccatct tcagtcttgc tgacagctaa 120
agagcaattg aaaccatttc cagcctaaac tacataaaag cagccgaacc aatgattaaa 180
gacctctaaag gccccataat catcattaaa tatgcccaca ctcatgttga ctttttattt 240
tatatacagg attaaaaatca acattaaatc attctattta catggccatc ggtgctgaaa 300
ttgagcattt taaatagtac agtaggctgg tatatcattag gaaatggact gcactggagg 360
caaatagaaa actaaagaaa ttagataggc tggaaatgct t                                     401

```

<210> 54

<211> 401

<212> DNA

<213> Homo sapiens

<400> 54

```

cccaacacaa tggataaaaa cacttatagt aaatggggac attcactata atgatctaag 60
aagctacaga ttgtcatagt tgttttctct ctttacaaaa ttgctccaga tctggaatgc 120
cagtttgacc tttgtcttct ataatttttc ctttttttcc cctctttgaa tctctgtata 180
tttgattctt aactaaaatt gttctcttaa atattctgaa tctctggtaat taaaagtttg 240
gggtattttt ctttacctcc aaggaaagaa ctactagcta caaaaaatat ttggaataaa 300
gcattgtttt ggtataaggt acatatatttg gttgaagaca ccagactgaa gtaaacagct 360
gtgcaccaa tttattatag ttttgtaagt aacaatatgt a                                     401

```

<210> 55

<211> 933

<212> DNA

<213> Homo sapiens

<400> 55

```

tttactgcctt ggcaaaagtac cctgagcatc agcagagatg ccgagatgaa atcagggaac 60
tcttagggga tgggtcttct attacctggg aacacctgag ccagatgcct tacaccacga 120
tgtgcatcaa ggaattgcct cgcctctacg caccggtagt aaactatccc ggttactcga 180
caaaaccatc acctttccag atggacgctc cttacctgca ggaataactg tgtttatcaa 240
tatttgggct cttaccacca acccctattt ctgggaagac cctcaggtct ttaacccctt 300
ggatttctcc agggaaaatt ctgaaaaaat acatccctat gccttcatac cattctcagc 360
tggattaaag aactgcattg ggcagcattt tgccataatt gagtgtaaag tggcagttgc 420
atttaactctg ctccgtttca agctggctcc agaccactca agggcaccca gctgtcgtca 480
agttgcctca agtccaaagaa tgggaatccat gtgtttgcaa aaaaagtttg ctaattttaa 540
gtctcttttc tataagaatt aakgagacaa ttttcttacc aaaggaagaa caaaaggata 600
aatataatag aaaatatatg tatatgggtg tttgacaaat tatataactt aggatccttc 660
tgactgggtt tgacatccat taacagtaat ttttaattct ttgctgtatc tgggtgaacc 720

```

```

cacaacaaac cctgaaaaaa ctcaagctga gttccaatgc gaagggaaat gatrgggttg 780
ggtaactagt ggtagagtgg ctttcaagca tagtttgatc aaaactccac ccagtatctg 840
cattactttt atctctgcaa atatctgcgt gatagcttta ttctcagtta cttttcccca 900
taataaaaaa tatctgccaa aaaaaaaaaa aaa
793

```

```

<210> 56
<211> 480
<212> DNA
<213> Homo sapiens

```

```

<400> 56
ggctttgaag catttttgc tgtgctccct gatcttcagg tcaccacccat gaagtcttta 60
gcagtcctgg tactcttggg agtttccatc tttctggctc ctgccagaa tcgcacaaca 120
gctgctccag ctgacacgta tccagctact ggtcctgctg atgatgaagc cccctgatgc 180
gaaaaccactg ctgctgcaac cactgcgacc actgctgctc ctaccactgc aaccaccgct 240
gcttctacca ctgctctgaa agacattcca gtttaccca aatgggttgg ggaatctccg 300
aatggtagag tgtgtccctg agatggaatc agcttgagtc ttctgcaatt ggtcacaact 360
attcatgctt cctgtgattt catccaacta cttaccttgc ctacgatatc ccccttatct 420
ctaactcagt tattttcttt caaataaaaa ataactatga gcaacaaaaa aaaaaaaaaa 480

```

```

<210> 57
<211> 798
<212> DNA
<213> Homo sapiens

```

```

<400> 57
agcctacctg gaaagccaac cagtcctcat aatggacaag atccaccagc tcctcctgtg 60
gactaaacttt gtgatatggg aagtgaatat agttaaacac ttgcacgacc aaacgaacga 120
agatgaccag agtactctta accccttaga actgttttcc cttttgtatc tgcaatatgg 180
gatgggtattg ttttcatgag ctcttagaaa ttccacttgc aagtttattt ttgcttcctg 240
tgttactgcc attcctattt acagtatatt tgagtgaatg atttatattt taataaagta 300
catggggctt ttttggttgt cctaaactta caaacattcc actcattctg tttgtaactg 360
tgattataat ttttgtgata atttctggcc tgattgaagg aaatttgaga ggtctgcatt 420
tatatatatt aaatagattt gataggtttt taattgtctt ttttccataa ggtatttata 480
aagttatttg gggttgtctg ggatttgtgt aaagaaaatt agaaccctgc tgtatttaca 540
tttacccttg tagtttattt gtggatggca gttttctgta gttttgggga ctgtggttagc 600
tcttggattg ttttgcaaat tacagctgaa atctgtgtca tgatttaaac tggcttatgt 660
ggctagaata ggaagagaga aaaaatgaaa tggttgttta ctaattttat actcccat 720
aaaattttta atgttaagaa aaccttaaat aaacatgatt gatcaatatg gaaaaaaaaa 780
aaaaaaaaa aaaaaaaa
798

```

```

<210> 58
<211> 280
<212> DNA
<213> Homo sapiens

```

```

<400> 58
ggggcagctc ctgacccctc acagccacct ggtcagccac cagctggggg aacgagggtg 60
ggggcagctc ctgacccctc gcttgcctcc gccactgcct tggtgcttgg tgatccaagt 120
ccctgcctcg gtccccaca aggaactcca tccaggccct ctctgcctcg ccccttgtca 180
tggacatagg tctgaggaaa gggctcatgc ccttatttta tgggaacctt ttcattctaa 240
cagaataaac cggaaggaa accagaaaaa aaaaaaaaaa
280

```

```

<210> 59
<211> 382

```

<212> DNA

<213> Homo sapiens.

<400> 59

```

aggcgggagc agaagctaaa gccaaagccc aagagagtgg cagtgccagc actggtgcca 60
gtaccagtag caataacagt gccagtgcc agtgccagcac cagtgtgtgc ttcagtgctg 120
gtgccagcct gacgcgccact ctccactttg ggctctctcg tggccttggg ggagctggtg 180
ccagcaccag tggcagctct ggtgcctgtg gttctctcta caagtggagt tttagatatt 240
gttaatcctg ccagctcttc tcttcaagcc aggggtgcac ctccagaaac tactcaacac 300
agcactctag gcagccacta tcaatcaatt gaagtgtgaca ctctgcatta aatctatttg 360
ccattaaaaa aaaaaaaaaa aa 382

```

<210> 60

<211> 602

<212> DNA

<213> Homo sapiens

<400> 60

```

tgaagagccg cgcgggtggag ctgctgccgc atgggaactgc caaccttgcc aagctgcagc 60
ttgtggttga gaatagtggc cagcgggtga tccacttgcc gggtcagtg gagaaagcacc 120
gggtcccatc ctctgtagta ccgccaactcc gaaagctgca ggattgcaga gacctggaat 180
ctctctcagc gctcggcagag atccaaagac tgcaccagag ttccggggcg gctctggaag 240
aggcccgacg gaaggaggag gtctataaag agctgatgtc agagctggag actctgcaca 300
ggagtgtgtc ccggctggcc tacaccagc gcactcctgga gatcgtggcg aacatccgga 360
agcagaagga agagatcacc aagatcttgt ctgatacgaa ggagcttcag aaggaataca 420
actccctatc tgggaagctg gacccggcgt ttgcggtgac tgatgagctt gtgttcaagg 480
atgccaaaga ggacgatgct gttcgggaag cctataagta tctagctgct ctgcacagga 540
actgcagcca gctcatccag accatcgagg acacaggcac catcatcgcg gaggttcgag 600
ac 602

```

<210> 61

<211> 1368

<212> DNA

<213> Homo sapiens

<400> 61

```

ccagtgagcg cgcgtaatac gactcactat agggcgaaat gggtaccggg cccccctcg 60
agcggccgcc cttttttttt tttttttatt gatcagaatt caggctttat tattgagcaa 120
tgaaaaacgc taaaaactaa ttccaagcat gtgtagttaa agtttgcaga gtgggataatt 180
gttcacaaaa cacattcaat gtttaaacac tatttatgtt aagaacaaaa tatattttaa 240
attgtttgct tctaaaaaag ccatttccct ccaagcttaa actttgtaat ttgatattaa 300
gcaatgaagt tattttgtac aatctagtta aacaagcaga atagcactag gcagaataaa 360
aaattgcaca gacgtatgca attttccaag atagcattct taaattcag ttctcagctt 420
ccaaagattg gtgtcccata atagacttaa acatataatg atggcctaaa aaaaataagta 480
tacgaaaaatg taaaaaagga aatgtaagtc cactctcaat ctcataaaag gtgagagtaa 540
ggatgctaaa gcaaaaataaa tgtagggtct tttttctgtg ttccggttat catgcaactc 600
gcttctttga tatgccttag ggttaccocat ttaagttaga ggttgaatg caatggtggg 660
aatgaaaaat gatcaaatat acaccttgct atttcatttc aaattgcggg ctggaaactt 720
ccaaaaaaag ggtaggcatg aagaaaaaaa aaatcmaatc agaacctctt cagggggttg 780
kgktctgata tggcagacac gatcaaatc ccaccaggag atggagcaat tcaaaataag 840
ggtaatgggc tgacaaggta ttattgccag catgggacag aatgagcaac aggcgtgaaa 900
gtttttggtat tatatagcac cttagagtctc tgatgtaggg aattttgtgt agtcaaacat 960
acgctaaact tccaaggaaa aatctttcag gttagcctaag ctgcttttc tagagtgatg 1020
agtgctattg ctactgtgat ttttgaaaa caaactgggt ttgtacaagt gagaagact 1080
agagagaaaag attttagtct gtttagcaga agccatttta tctcgctgca catggatcaa 1140

```

```

tattttctgat cccctataacc ccaggaagg gcaaaatccca aagaaatgtg ttagcaaaat 1200
tggtctgatgc tatcatattg ctatggacat tgatcttgcc caacacaatg gaattccacc 1260
acactggatc agtggatcca ctagtcttag agcggcggcg caccgcggtg gagctccacc 1320
ttttgttccc tttagtggag gtttaattgcg cgcttgggct aatcatnn 1368

```

<210> 62

<211> 924

<212> DNA

<213> Homo sapiens

<400> 62

```

caaaagnaca ggaacagctt gnaaagtact gncatnccn cctgcaggga ccagcccttt 60
gcctccaaaa gcaataggaa atttaaaaga tttnccactga gaaggggnc cagcttnart 120
tntnaatgtn tcargnanar tnccctncaa atgncrncn cactnaetnr gnatttgggt 180
tnccgnrtnc mgnactatnt caggtttgaa aaactggatc tgccacttat cagttatgtg 240
accttaaaaga actccgctaa ttcttcagag cctcagtttc ctgtctata agttgggagt 300
aatattaata ctatcttttt tccaaggatt gatgtgaaca ttaatgaggt gaaatgacag 360
atgtgtatca tgggtcctaa taacatcca aatatatga cttactatgt tcattattat 420
tacttggttg aagcttaaga cctcacata gaatcccatc cagcccacca cacagaggtc 480
tgagttttct agtttggaa agctattaaa taacaacttc tagtgtcaat tcataactgt 540
ttatgggtcaa gtaactgggc tcagcatttt acattcattg tccttttaag ttctagcaat 600
tgaagcagg aaactatgatt atattgacta cataaatgaa gaaattggag ctcatagaca 660
ttaagtaatt ctcccagggt cacacagcta gaactggcaa agcctgggag tgatccatga 720
tcttcagaca ttgaagaatc ataatgttaa ataatgtcaa ggccttttcc tcagaaagag 780
tcctgggtgc tgacccaacc cactagcact tgttctctac aggggaacat ctggtggcct 840
gggaactcact gcacgtcgca agagatgttg cttctgatga attattgttc ctgctcagtg 900
tgtgaaggga aaaaaaaaaa aaaa 924

```

<210> 63

<211> 1079

<212> DNA

<213> Homo sapiens

<400> 63

```

agtcaccaaga actcaataat ctcttatggt ttcttttgaa gacttatttt aaatattaac 60
tatttcgggt cctgaaatgga aaaatataaa cattagctca gagacaatgg ggtacctgtt 120
tggaatccag ctggcagcta taagcacctg tgaaaaactc gacaggcttt ttgccctttt 180
tattaaatgg cctcacatcc tgaatgcagg aatgtgttgc ttttaaaaaa cattaacttt 240
taatgttgaa ttctgaaaac acaaccataa atcatagttg gtttttctgt gacaatgac 300
agtcacatta ttctctccac agcaaaacta cctttccaga aggtggaaat tgtatttgca 360
acaatcagg caaaacccac actgaaaaag cattttacaa tattatatct aagttgcaca 420
gaagaccaca gtgatcacta ggaatctac cacagtcagg ttttctaat ccaagaaggt 480
caaaactctg ggaataatgt gtccctcttc tgctgctgct ctgaaaaata ttcatcaaa 540
acgaagttta caagcagcag ttattccaag attagagttc atttggat ccactgtata 600
ctggcaatgt ttaggtttgc ccaaaaactc ccagacatcc acaatgtgtg tgggtaaaac 660
accacatctg gtaacctctc gatcccttag atttgcatt ctgcaaaa taactgtagc 720
tgactctgga gctccttgca ttttctttaa aaccattttt aactgatca ttctgtccg 780
agcatgccct ctggtgctct ccaaatggga tgcataagg caaagctcat ttctgacac 840
attcacatgc acacataaaa ggtttctcat cattttggtt cttggaaaaa gaataatccc 900
ttggcttttt aatttcaact ttgatttctt caacattata gctgtgaaat atccttcttc 960
atgacctgta ataactcat aattacttga tctctcttct aggtagctat aatatggggg 1020
aataactctc tgtagaataa tcacatctgg gctgtacaaa gctaaagtag acacaccc 1079

```

<210> 64

<211> 1001

<212> DNA

<213> Homo sapiens

<400> 64

```

gaatgtgcaa cgatcaagtc aggggtatctg tggatccac cactttgagc attatcgat 60
tctatatgtc aggaacattt caagttatct gtcttagcaa ggaataataa aatacttata 120
gttaactatg gcctatctac agtgcaacta aaaactagat tttattcctt tccacctgtg 180
ggtttgattt catttaccac cctcttttca ttccctttct caccacaca ctgtgccggg 240
cctcaggcat atactattct actgtctgtc tctgtaagga ttatcatttt agcttccaca 300
tatgagagaa tgcattgcaa gtttttcttt ccatgtctgg ctattttcac ttaacataat 360
gacctcgctt tccatccatg ttattttatat taccataatg tgttcataaa tatatatata 420
cacatatata ccacattgca ttgttccaat tattcattga cggaacttgg ttaatgttat 480
actcgttgcta ttgtggtaga tgcgtgcaata aaacagcgaag tgggggataa atttgaagag 540
ttttttggtt gatgttcttc caaattttaa gattgttttg tctatgtttg tgaatatggc 600
gttagtattt tcatagagat tgcattgaaat ctgtagattg ctttgggttaa gtatggttat 660
tttgatggtt ttaatttttt cattccatga agatgagatg tctttccatt gtttgtgtcc 720
tctacatttt .ctttcatcaa agttttgttg tatttttgaa gtatagtat ttcaccttat 780
agatcaagtg tattccctaa atattttatt ttgtagacta ttgtagata aattgccttc 840
tgtattctt ttccctctaa ttcatatta gtgtatgtaa atgttatgga tttttatttg 900
tgggttttta atcaaaaact gtattaaact tagagttttt tgtggagttt ttaagttttt 960
ctagatataa gatcatgaca tctacaaaaa aaaaaaaaaa a 1001

```

<210> 65

<211> 575

<212> DNA

<213> Homo sapiens

<400> 65

```

acttgatata aaaaggatat ccataatgaa tattttatc tgcaccttt acattagcca 60
ctaaatcagt tattgtttga tgaagacctt tcacagaatc ctatggattg cagcatttca 120
cttggctact tcatacccat gccttaagaa ggggcagttt ctcaaaagca gaaacatgcc 180
gccagttctc aagttttctt cctaactcca ttggaatgta agggcagctg gccccaatg 240
tggggaggtc cgaacatttt ctgaattccc attttcttgt tcggcgctaa atgcagttt 300
ctgtcattac ttagattccc gatctttccc aaaggtgttg atttacaagg aggccagcta 360
atagccagaa atcatgaccc tgaagagag atgaatttc aagctgtgag ccaggcagga 420
gccccagtat ggcaaaaggtt ctgtgagaatc agccatttgg tacaataaag atttttaag 480
cttttatggt ataccatgga gccatagaaa ggctatggat tgtttaagaa ctatttttaa 540
gtgttccaga cccaaaaagg aaaaaaaaaa aaaa 575

```

<210> 66

<211> 831

<212> DNA

<213> Homo sapiens

<400> 66

```

atcgggtctc ttctgctaaa cagccacatt gaaatgggtt aaaagcaagt cagatcagggt 60
gatttgtaaa attgtattta tctgtacatg tatgggtttt taattcccac caagaaagag 120
agaaaatttc tttttagtta aaaccaaatt tcacttttca aaatatcttc caacttattt 180
attcgtgttc actcaattgc ctatataat atatatatat gtgtgtgtgt gtgtgtgcgc 240
gtgagcgcac ctgtgtgtat gcgtgcgcac gtgtgtgtat gtgtattatc agacataggt 300
ttctaaactt tagatagaag aggaagcaaca tctatgccaa atactgtgca ttctacaatg 360
gtgctaaact cagacctaaa tgatactcca ttaattttaa aaaagagttt taaataatta 420
tctatgtgct tgtatttccc ttttgagtgc tgcacaacat gtttaacata tagtgtaaaa 480
cgaagtgaac caaccacgtg ttctaaagtc tagggattgt gctataatcc ctatttagtt 540
caaaataaac cagaattctt ccattgtgaa tggaccaaac tcatattatt gttatgtaaa 600

```

```

tacagagttt taatgcagta tgacatccca caggggaaaa gaatgctctg agtgggtgac 660
tgttatcaaa tattttatag aatacaatga acggtgaaca gactggtaac ttggttgagt 720
tccatgaca gatttgagac ttgtcaatag caaatcatct ttgtatttaa attttgtac 780
tgatttgaaa aacatcatta aatatcttta aaagtaaaaa aaaaaaaaaa a 831

```

<210> 67
 <211> 590
 <212> DNA
 <213> Homo sapiens

```

<400> 67
gtgctctgtg tattttttaa ctgcattaga cattgaatag taatttgcgt taagatacgc 60
ttaaaggctc ttgtgacca tgtttccctt tgtagcaata aaatgttttt tacgaaaaact 120
ttctccctgg attagcagtt taaatgaaac agagttcatc aatgaaatga gtatttaaaa 180
taaaaatttg ccttaatgta tcagttcagc tcacaagatg tttaagatga ttgagagac 240
ttgaattaaa gaaaaaaaaa ttctcaatca tattttttaa atataagact aaaattgttt 300
ttaaaacaca ttccaatag aagtgaagtt gaactgacct tatttatact ctttttaagt 360
ttgttccctt tccctgtgcc tgtgtcaaat ctccaagctc tgctgaaat acatttgata 420
caaaagtctt tgtagttgtg tttagttctt tgtcatgtct gtttttggct gaagaaccaa 480
gaagcagact ttctttttaa aagaattatt tctctttcaa atatttctat ccttttttaa 540
aaattccttt ttatggctta tatacctaca tatttaaaaa aaaaaaaaaa 590

```

<210> 68
 <211> 301
 <212> DNA
 <213> Homo sapiens

```

<400> 68
ttgtgttggg gttccctttt ccggtcggcg tggctcttgc agtggagtg ccgctgtgcc 60
cgggcctgca ccatgagcgt ccggtccttc atcgacatca gtgaagaaga tcaggctgct 120
gagcttcgtg cttatctgaa atctaaagga gctgagattt cagaagagaa ctcggaaggt 180
ggacttcatg ttgatttagc tcaaatattt gaagcctgtg atgtgtgtct gaaggagat 240
gataaagatg ttgaagtgt gatgaacagt ggggnatcct actcttgatc cggaanccna 300
c 301

```

<210> 69
 <211> 301
 <212> DNA
 <213> Homo sapiens

```

<400> 69
tctatgagca tgccaaggct ctgtgggagg atgaaggagt gcgtgcctgc tacgaacgct 60
ccaacgagta ccagctgatt gactgtgccc agtacttctt ggacaagatc gacgtgatca 120
agcaggctga ctatgtgccg agcgatcagg acctgcttgc ctgccgtgtc ctgacttctg 180
gaatctttga gaccaagttc caggtggacn aagtcaactt ccacatgntt gacgtgggtg 240
gccagcgaga tgaacgccgc aagtggatcc agtgcttcaa cgatgtgact gccatcatct 300
t 301

```

<210> 70
 <211> 201
 <212> DNA
 <213> Homo sapiens

```

<400> 70
gcggctcttc ctccggcagc ggaagcggcg cggcggctcg agaagtggcc taaaacttcg 60

```

gcgttgggtg aaagaaaaat gccccaacca agcagactgc tcgtaagtcc accgggtggga 120
 aagcccccgc caaacagctg gccacgaaag ccgcccaggaa aagcgcctccc tctaccggcg 180
 gggtagagaa gccctcatgc t 201

<210> 71
 <211> 301
 <212> DNA
 <213> Homo sapiens

<400> 71
 gccggggtag tcgccgncgc cggccgcccgt gcagccactg caggcaccgc tgccgcccgc 60
 tgagtggtgg gcttaggaag gaagagggtca tctcgctcgg agcttcgctc ggaagggtct 120
 ttgtccctg cagccctccc acgggaatga caatggataa aagtgcagctg gtacanaaag 180
 ccaaacctgc tgagcaggct gagcgatatg atgatatggc tgagcccatg aaggcagtc 240
 cagaacaggg gcatagaact ttcaacgaag agagaaatct gctctctggt gcctacaaga 300
 a 301

<210> 72
 <211> 251
 <212> DNA
 <213> Homo sapiens

<400> 72
 cttggggggt gttgggggag agactgtggg cctggaaata aaacttgtct cctctaccac 60
 caccctgtac cctagcctgc acctgtccac atctctgcaa agttcagctt ccttccccag 120
 gtctctgtgc actctgtctt ggatgctctg gggagctcat ggggtgggga gtctccacac 180
 gaggggaggt caggggactg gttgggccag ggatgaatat ttgagggata aaaatttgtt 240
 aagagccaan g 251

<210> 73
 <211> 913
 <212> DNA
 <213> Homo sapiens

<400> 73
 tttttttttt tttttcccat gccctctttt tatttacagt gataccaaaac catccacttg 60
 caaattcttt ggtctcccat cagctggagt taagttagta ctgtgtatct ttgagatcat 120
 gtatttggct ccactttggg ggatacaaga aaggaaggca cgaacagctg aaaaagaagg 180
 gtatocaccc gtcacagctg gaatccagca ggaacctctg agcatgccac agctgaacac 240
 tcaaaaggag aaagaaggac agctgctctt cattttttt gaaagcaaat tcatttgaaa 300
 gtgcataaat ggtcatcata agtcaaacgt atcaataga ccttcaacct aggaacaaca 360
 attttttttt tctatttaat aatacaccac actgaaatta ttggccaatg aatcccaagg 420
 atttggtaaca atagttacaa ttctgtattg ctttctctct tctttctctc agacaaacac 480
 caaatataat gcaggtgaaa gagatgaacc acgactagag gctgacttag aaatttatgc 540
 tgactcgatc taataaaaaat tatgttgggt aatgttaact tatctataat agagcatttt 600
 ggggaatgct ttcaaaaagg gtcagatcac agtcatacag ctgaaaaagt cccctgaaaaa 660
 aagaatggtt aagaagtata ataacctttt caaaacccac aatgcagctt agttttctct 720
 tattttattg ttggtcatgaa gactatcccc atttctccat aaaaatctct cttccatactg 780
 ctgcatcatg gcacaaaaga ctctaagtc caccagacag aaggaccaga gtttctgtatt 840
 ataaacaatg atgctgggta atgttttaaa gagaacattg gatatggatg gtcagcccaa 900
 cacaatgaa ttc 913

<210> 74
 <211> 351
 <212> DNA
 <213> Homo sapiens

<400> 74
 tgtgcnacagg ggaagggtgg gcngtgagga ngatgacaga aaggctggaa ggaanggggg 60
 tgggtttgaa ggcacnggcc aaggggncct cagggtccgnt tctgnnaagg gacagccttg 120
 aggaaggagn catggcaagc catagctagg ccaccaatca gattaagaaa nncatgagaaa 180
 nctagctgac catcactgtt ggtgnccagt ttcccaacac aatggaatnc caccacactg 240
 gactagnnga hccactagtt cttagagcgg cgccaccggc gtggaacccc aacttttggc 300
 ccttagnga gggtaattg cgcgcttggc ntaatcatgg tcataagctg t 351

<210> 75
 <211> 251
 <212> DNA
 <213> Homo sapiens

<400> 75
 tactgacact tctttgaaaa gcattcccaa aatgctctat tttagataga ttaacattaa 60
 ccaacataat tttttttaga tcgagtcagc ataaatttct aagtcagcct ctagtctggg 120
 ttactctctt tcacctgcat tttatttggg gtttgtctga agaaaggaaa gaggaaagca 180
 aatcagaatt gtactatttg taccaaatct ttgggattca ttggcaaaata atttcagctg 240
 ggtgtattat t 251

<210> 76
 <211> 251
 <212> DNA
 <213> Homo sapiens

<400> 76
 tatttaataa tacaccacac tgaaattatt tgccaatgaa tcccaaagat ttggtacaaa 60
 tagtacaatt cgtatttggc ttctcttttc ctttcttcag acaaacacca aataaaatgc 120
 aggtgaaaga gatgaaccac gactagaggg tgacttagaa atttatgtct actcgatcta 180
 aaaaaaatta tgggtggtta tgtaaatcta tctaaaatag agcatttttg gaatgctttt 240
 caaagaaggt c

<210> 77
 <211> 351
 <212> DNA
 <213> Homo sapiens

<400> 77
 actcaccgtg ctgtgtgctg tgtgcctgct gcctggcagc ctggccctgc cgctgctcag 60
 gaggcgggag gcatgagtg gctacagtg gaacaggctc aggcatactt caagagannn 120
 tatctctatg actcagaaac aaaaaatgcc aacagtttag aagccaaact caaggagatg 180
 caaaaattct ttggcctacc tataactgga atgttaaact cccgcgtcat agaaataatg 240
 cagaagccca gatgtggagt gccagatgtt gcagaatact cactatttcc aaatagccca 300
 aaatggactt ccaagtgtg cactacagg atcgtatcat atactcgaga c 351

<210> 78
 <211> 1592
 <212> DNA
 <213> Homo sapiens

<400> 78
 gaattccatt gtgttggggc cctgggggag ggggggaggg gccaccacg gccttatttc 60
 cgcgagcgcc ggcactgcc gctccgagcc cgtgtctgtc ggggtgccag ccaactttcc 120
 tgcgtccatg cagcccgccg ggcaacggct gcccgctccc tgggtccggc ccagggggccc 180
 gcgcgccacc gcccccgctc tcgocgtgct gctgtttgct gccccgggtg cggcgcccgcc 240
 ggggtccggg gaccocgagc accctgggca gcttcaggat gctggggctc cgcgcaggct 300
 cctgcagcag gcggcgcgcg cggcgcttca cttcttcaac ttccgggtcc gctcgccagc 360
 cgcgctcgga gtgctggccc aggtgcagga gggcgcgcg tggattaat caaaagaggg 420
 atgtaaagt cagctggtct tcagcacaga gcgctacaac ccagagtcct tacttcagga 480
 aggtgaggga cgtttgggga aatgttctgc tcgagtgttt ttcaagaatc agaaacccag 540
 accaactatc aatgtaact gtacacggct catcgagaaa aagaaaagac aacaagagga 600
 aaagctcttac gtgatgtggg aatgacaact acagggtgtca cactactact tggcacagct 660
 tgataatcat ggacatatg atccctctct gagactcatc tgggatttgg ctttccctgg 720
 cactagtgtg aggcagtgga aaactaatga tgatacaatt gattttgatt atactgttct 840
 acttcattga ttatcaacac aggaataaat tcctgtctgc attcacttgg tctgggtacc 900
 tggcaaacct cttaaagtga agtaccactg tcaagagcta cagacaccag aagaagcctc 960
 cggaaactga gaagcagcag ctgtagtacc aacagagctt agtaatttct aaaaagaaaa 1020
 aatgatcttt ttccgagctc taacaagtg actatactag cataaatcat tcttctagta 1080
 aaacagctaa ggtatagaca ttctaataat ttgggaaaaa ctatgattac aagtaaaaaa 1140
 tcagaaaatgc aaagatgttg gttttttgtt tctcagctcg ctttagcttt taactctgga 1200
 agcgcatgca cactgaactc tgctcagtcg taaacagctc ccagcaggtt cctcaggggt 1260
 tcagccctaa aatgttaaac ctggataaac agtctatggt gcaccagagc cagcattttt 1320
 tttttaactg caaaaaatga ttgtctctac tctgaattta tatttctcat tcttttgaac 1380
 atactatagc taatatattt tatgttgcta aatgtcttct acttagcatc ttaataaaga 1440
 ataatactat ttcatgaaa gttaattata ggaaaaaaat taactgtttt aaaaagaaat 1500
 tgattatggt ttatgatttc aggcaagtat tcatttttaa cttgtctact actttttaat 1560
 aaatgtttac atttctaaaa aaaaaaaaaa aa 1592

<210> 79

<211> 401

<212> DNA

<213> Homo sapiens

<400> 79
 catactgtga attgtttctg actccttttc ttgacattca gttttcanaa ttccatctt 60
 tcttctgaa ctaagtgtct gttctcttga ctgcctgtcg ggccagcacc cgattgccag 120
 ccagaaacgt cacactgcc cagatggcca ggtacttcaa ggtctggaac atgttgagct 180
 gagtccagta gacatacatg agtcccagca tagcagcatg tcccagggtga aatataatcg 240
 tgctaggagc aaaaagtgaag ttggagacat tggccaccaat caggtccacc tagttctaga 300
 gcggcccgca ccgcggtgga gctccagctt ttgttccctt ttgtgaggt taattgcgag 360
 cttggcgtaa tcatggnatc agctgttttc tgtgtgaaat t 401

<210> 80

<211> 301

<212> DNA

<213> Homo sapiens

<400> 80
 aaaaatgaaa catctatttt agcagcaaga ggctgtgagg gatggggtag aaaaggcatc 60
 ctgagagagt tctagaccga cccaggtcct gtggcacact atacgggtga ggaaggggtg 120
 aagacagggc taagctctag gacgggtgaat ctgggggcta tttgtggatt tgttagaacc 180
 agacattctt ttggcctttt cctggcactg gtgttgccgg caggtgggga gaagtgaacc 240
 accagtcact gttcagtcac tgccaccaca gatcttcagc agaattctcc ggtaatcccc 300
 t 301

<210> 81
 <211> 301
 <212> DNA
 <213> Homo sapiens

<400> 81
 tagccagggtt gctcaagcta attttattct ttcccaacag gatccatttg gaaaatatca 60
 agcctttaga atgtggcagc aagagaaagc ggactacgca ggaacgggga gtttgggaga 120
 agctctcctg gtgttgactt agggatgaag gctccaggct gctgccagaa atggagtcac 180
 cagcagaaga actgntttct ctgataagga tgtccacca ttttcaagct gttcgttaaa 240
 gttacacagg tcctctctgc agcagtaagt accgttagct cattttcct caagcgggtt 300
 t 301

<210> 82
 <211> 201
 <212> DNA
 <213> Homo sapiens

<400> 82
 tcaacagaca aaaaaagttt attgaataca aaactcaaag gcatcaacag tcctgggccc 60
 aagagatcca tggcaggaag tcaagagttc tgcttcaggc tcggtctggg cagccctgga 120
 agaagtcatt gcacatgaca gtgatgagtg ccaggaaaac agcatactcc tggaaaagtc 180
 acctgctggn cactgnttca t 201

<210> 83
 <211> 251
 <212> DNA
 <213> Homo sapiens

<400> 83
 gtaaggagca tactgtgccc attttattata gaatgcagtt aaaaaaata ttttgagggt 60
 agcctctcca gtttaaaagc acttaacaag aaacacttgg acagcgatgc aatggtctct 120
 cccaaaccgg ctccctctta ccaagtaccg taaacagggt ttgagaacgt tcaatcaatt 180
 tcttgatatg aacaatcaaa gcatttaagt caaacatatt tgctctctca anaataaac 240
 cattttccaa a 251

<210> 84
 <211> 301
 <212> DNA
 <213> Homo sapiens

<400> 84
 agtttataat gttttactat gatttagggc ttttttttca aagaacaaaa attataagca 60
 taaaaactca ggtatcagaa agactcaaaa ggctgttttt cactttgttc agattttgtt 120
 tccaggcatt aagtgtgtca tacagttgtt gccactgctg ttttccaaat gtccgatgtg 180
 tgctatgact gacaactact tttctctggg tctgatcaat tttgcagtan accattttag 240
 ttcttacggc gtcnataaca aatgcttcaa catcatcagc tccaatctga agtcttgctg 300
 c 301

<210> 85
 <211> 201
 <212> DNA
 <213> Homo sapiens

<400> 85
 tattttgtgta tgtaacattt attgacatct acccactgca agtatagatg aataagacac 60
 agtcacaccca taaaggagtt tatectttaa agggagtgaag gacattcaaa aaccaactgc 120
 aataaaaaag ggtgacataa ttgctaaatg gaggaggagga acagtgctta tcaattcttg 180
 attgggccac aatgatatac c 201

<210> 86
 <211> 301
 <212> DNA
 <213> Homo sapiens

<400> 86
 tttataaaat attttattta cagtagagct ttacaaaaat agtcttaaat taatacaaat 60
 ccccttttga atataactta tatgactatc ttctcaaaaa cgtgacattc gattataaca 120
 cataaactac atttatagtt gtttaagtcac ctgttagtat aaatatgttt tcatcttttt 180
 tttgtaataa ggtacatatt aataacaatg aacaatggac acaaatcttt attttgntat 240
 tcttccaatg taaaattcat ctctggccaa acaaaaatta accaaagaaa agtaaaacaa 300
 t 301

<210> 87
 <211> 351
 <212> DNA
 <213> Homo sapiens

<400> 87
 aaaaagatt taagatcata aatagggtcat tgtgtgcaca acacatttca gaattttaaa 60
 aaaaacaaca ttttggcttt ctaagaaaaa gacttttaaa aaaaatcaat tccctcatca 120
 ctgaaggagc ttgtacattt ttaactcttc agtctcctaa ggcacagtat ttaatcagaa 180
 tgccaatatt accaccctgc tgtagcanga ataaagaagc aagggattaa cacttaaaaa 240
 aacngccaaa ttccctgaac aatcatttgg catttttaaa aagggataaa aaaaacnggt 300
 aaggggggga gcattttaag taaagaangg ccaaggggtg tatgccngga c 351

<210> 88
 <211> 301
 <212> DNA
 <213> Homo sapiens

<400> 88
 gttttaggtc tttaccaatt tgattggttt atcaacaggg catgagggtt aaatatatct 60
 ttggggaag gtaaatgcaa atttgacttc atagggtcgc ggctcctca ctccctgtgca 120
 tttcttggtg gaagcacaca gtttaattaac tcaagtgtgg cgttagcgat gctttttcat 180
 gmgctcatt atccacttgg tgaacttgc cacttgaatg naaactcctg ggtcattggg 240
 ntggccgcga gggaaaggtc cccaagacac caaaccttgc aggttacctn tgacaccaa 300
 c 301

<210> 89
 <211> 591
 <212> DNA
 <213> Homo sapiens

<400> 89
 tttttttttt tttttttt aatcaaatga ttcaaaaaca ccatcattct gtcaatgcc 60
 aagcacccag ctggtcctct cccacatagt cacactctcc tcagcctctc ccccaacctt 120
 gctctccctc ctcccctgcc ctaggccagg gacagagctt agggaggacc tggggcagag 180
 ctggaggcag gaagagagca ctggacagac agctatggtt tggattgggg aagagattag 240

gaagtaggtt	cttaaagacc	ctttttttagt	accagatatc	cagccatatt	cccagctcca	300
ttattcaaat	catttcccat	agccagagctc	ctctctgttc	tcceccctact	accaatctct	360
tggctcttac	acaattttta	tccttcaaat	attcatccct	ggcccaacca	gtccctctgag	420
ctctccctctg	gtggagactc	ctccaccat	gagctcccca	gagcatccaa	gacagagtgc	480
acagagacct	ggggaaggaa	gctgaacttt	gcagagatgt	ggacagggtgc	aggctagggt	540
acagggtggg	ggtagaggag	acaagtttta	tttccaggcc	cacagtctct	c	591

<210> 90

<211> 1996

<212> DNA

<213> Homo sapiens

<400> 90

tttttttttt	ttttttatca	aatgaatact	ttattagaga	cataacacgt	ataaaataaa	60
ttttttttca	tcattggagt	accagatatt	aaaaccaacc	aacactttct	cattttttaca	120
gtcaagacat	gttaaattct	taaatgccat	aatttttggt	caactgtctt	gtcatccaac	180
tcacagtcct	agaattctgt	taagctacaa	atctcaagt	tcacagatgt	gtcttagggt	240
tggttgttaa	caatctagaa	gcaatctggt	tacaaaagtg	ccaccaaaagc	atttttaaga	300
aaccaattta	atgccaccaa	acataagcct	gctatacctg	ggaaacaaaa	aatctcacac	360
ctaaattcta	gcagagtaaa	cgatttccaac	tagaatgtac	tgtatatacca	tatggcacat	420
ttatgacttt	gtaaatatga	attcataata	cagggtttagg	tgtgtgggtat	ggagctagga	480
aaacccaaagt	agtaggatatt	tatagaaaag	atctgatgtt	aagtataaag	tcatatgcct	540
gatttctcca	aaacctttgt	ttttcctcat	gtctctctgtc	tttatatttt	tatacaaaac	600
caagatctaa	cagggttcttt	tctagaggat	tattagataa	gtaacacttg	atcattaaagc	660
acggatcatc	ccactcatct	atggttggtc	tatgttccat	gaactctaat	agcccaactt	720
atacatggga	ctccaagggg	atgcttcagc	cagaaagtaa	agggctgaaa	aagtagaaca	780
atacaaaagc	cctcgtgtgg	tgggaactgt	ggcctcactc	ttacttgtcc	ttccattcaa	840
aacagtttgg	cacctttcca	tgacgaggat	ctctacaggt	aggttaaaat	acttttctgt	900
gctatttcagc	cagaaaatgt	ttttgtgctg	gatatgattt	taaaacagat	tttgtctgtc	960
accagtgcaa	aaacattaca	gatgtctggg	ctaatacaaa	aacacataag	aatctacaac	1020
ttttatatta	atactctatt	caaatttaac	tcaaagtaat	gcataataat	tagaagtaaa	1080
aacttaattc	tcttgagagc	tctatttgga	aaagcttcac	atatccacac	acaaatattg	1140
gtatatattc	gcacagggca	aaacactgta	ttctgaagca	taataaaact	caaagtaaga	1200
catcagtacg	tagataaccg	ttccagtatt	ggtaaatggg	ctctggggat	cccattttaa	1260
gcactctcag	ataggagatt	tgctcagttg	ttagactatt	attagtttga	ttaagcaact	1320
gaagttttact	tcataaatac	ctttttccta	tatccaggac	tctgcctctg	aaattttata	1380
cattctccca	aaagtaagta	ttctccaaag	gtaagatttt	gactattaac	acaaggacaa	1440
tgtgattatt	gcataatgac	actaaatatt	atgtggcttt	tctgttaggt	ttataagttt	1500
tcaatgattc	gttcaaagaa	atgcagatca	tataataact	aggttttaca	ccagtggttg	1560
acaaactatg	gcccaagggc	taaaccaccg	ctcccttgtt	ttttataaat	aagttttatt	1620
agacataaac	acaactattc	atttctgtat	tgtgtatagc	tgctttcacg	ctataactagc	1680
agaactgaat	agttgtgaca	gagactgtat	ggaccgtgaa	gcataaatat	taccactctg	1740
gcccatctta	aaaaaagtgt	gcacaaattc	ggtttacact	aaaataatga	gttttagtggg	1800
aagctatttt	gaaatgtgtt	tttttttagg	gctgtaatta	ccaaattaaa	ttaaaggtcca	1860
ggtgactcag	caaccacaaca	aaagggatcc	taatttttta	tgaacaaat	atttgtattt	1920
tatggacata	aaaggaaact	ttcagaaaga	aaaggaggaa	aataaaagggg	gaaagggacc	1980
caacacaaatg	gaattc					1996

<210> 91

<211> 911

<212> DNA

<213> Homo sapiens

<400> 91

gccccttttt	tttttttttt	cttggttaaa	aaaattgttt	tcattttaat	gatctgggtt	60
------------	------------	------------	------------	------------	------------	----

agtaacaaac	aaatgtacaa	aattgtcttt	cacatttcca	tacattgtgt	tatggaccaa	120
atgaaaaacg	tggactacaa	atgcagggtt	ctttatatcc	ttaacttcaa	ttattgtcac	180
ttataaataa	aggtgatttg	ctaacacatg	catttgtgaa	cacagatgcc	aaaaattata	240
catgtgaagt	aatgcacaac	caagagtata	cactgttcat	ttgtgcagtt	atggctcaaa	300
tgcgactgac	acagaagcag	ttatcctggg	atatcttcat	ctatgatgaa	agcactcttg	360
agaaatagat	tgaatacag	tttaaaacga	aaattgtatt	ctacaataac	aataaaattt	420
gcaacttgca	catctgaagc	aacatttgag	aaagctgctt	caataaccct	gctgttatat	480
tgggtttata	ggatatcttc	caaagtcag	gggtgggata	tagctgcttt	aaagaaaaata	540
aatatgtata	ttaaaaggaa	aatcacactt	taaaaatgtg	agggaaagctt	tgaaaaacagt	600
cttaatgcac	gagtcacatc	acatattttc	aagtttttga	aacagaaaaga	agtttagaat	660
tttcaaaagta	atctgaaaac	tttctaagcc	attttaaaat	aagatttttt	tccccacttt	720
tccaatgttt	ctattttgat	agtgtaatac	agaaaatgggc	agtttctagt	gtcaacttaa	780
ctgtgctaet	tcataagtca	ttatacattt	atgacttaag	agttcaataa	agtggaaatt	840
gggttataat	gaaaatgaca	agggggcccc	ttcagcagcc	actcatctga	actagtaate	900
ccaacacaat	g					911

<210> 92

<211> 1710

<212> DNA

<213> Homo sapiens

<400> 92

tttttttttt	tttttaactt	tttagcagtg	ttatttttgt	taaaagaaac	caattgaatt	60
gaagggtcaag	acaccttctg	attgcacaga	tttaacaaga	aagtattact	tatttcaact	120
ttacaagaaga	tcttatttgt	ttaaaaagat	ccatactatt	gataaaagttc	accatgaaca	180
tatatgtaat	aaggagacta	aaatattcat	tttaccatct	tacaacatgt	atttcatatt	240
tctaatacaac	cacaaatcat	ataggaaaat	atttaggtcc	atgaaaaagt	ttcaaaacat	300
taaaaaatta	aagttttgaa	acaaatcaca	tgtgaagact	cattaaaaaa	taacattgac	360
aaataaatag	ttaatcagct	ttacttatta	gctgctgcc	tgcatttctg	gcattccatt	420
ccaagcgagg	gtcagcatgc	agggtataat	ttcatactat	gcgaccgtga	agagctacag	480
ggcttatttt	tgaagtga	tgtcacaggg	tctttcattc	tctttcaaa	gaagatcact	540
catggctgct	aaactgttcc	catgaagagt	accaaaaaag	cacctttctg	aaatgttact	600
gtgaagattc	atgacaacat	atttttttta	acctgttttg	aaggagtttt	gtttaggaga	660
ggggatgggc	cagtagatgg	agggtatctg	agaagccctt	ttctgtttta	aaataaatg	720
attcactgat	gtttatagta	tcaacagtct	tttaagaaca	atgaggaatt	aaaactacag	780
gatactgtgga	attttaaagc	aaattgcatt	catggatata	cctacatctt	gaaaaacttg	840
aaaaggaaaa	actattccca	aagaaggtcc	tgatacttaa	gacagcttgc	tgggtttgat	900
caaaagcagaa	agcatatact	ttcaagtgag	aaaacagcag	tggcaggctt	gagcttcca	960
agcaatcaaa	tctgtaaacg	agatggttac	tagtaagctc	agttatggga	gctcaggttc	1020
taactcatcag	tgtgcttctg	ggatttgctg	gctcttttcc	gctctctctg	atgctggact	1080
ggcttctcag	gtgacatgct	ctcaaaagtg	tgactggact	gtgtgtgctg	ccgggtgtac	1140
ctcttgcact	tgcaggcagt	gactactgtg	attttgtagg	tgcgtgtgct	gccactctgg	1200
cactgcagct	ggattctctg	ggtagcgggt	ttgtcattga	cacaccgcca	ctcctgggag	1260
ctcctctctg	tccagtaact	tgttccatag	ctctctccaa	tccagttagg	gagcactagg	1320
agggggcaag	actgcgcagc	acacaccagc	tccttcagag	ggctgatgct	ggtagactgg	1380
ccatcagaga	tgtatttggt	ggaaacagct	tcccggcgaac	ccacttgaac	ccgagtggtc	1440
cgatccagtc	cagtggttact	gaaatgcctg	cttccatttc	tggcttgatt	caactgtgct	1500
ttgtctgcgg	gggtgtctgg	aacaggttta	accacatgtg	aataaaggat	ttctgtggca	1560
tcatttttaa	aagccaaaca	gcttttcttt	aggatgcatg	caagggggaag	gagatagaaa	1620
tgaatggcag	gaggaagcat	ggtagtaga	ggatttgctt	gactgaagag	ctgggttaatt	1680
cttttgcctc	tgcccaaac	aatggaattc				1710

<210> 93

<211> 251

<212> DNA

<213> Homo sapiens

<400> 93

```

cccaccctac ccaaatatta gacaccaaca cagaaaagct agcaatggat tcccttctac 60
tttgtttaaat aataaagtta aatatttaaa tgcctgtgtc tctgtgatgg caacagaagg 120
accaacagggc cacatcctga' taaaaggtaa gaggggggtg gatcagcaaa aagacagtgc 180
tgtgggctga ggggacctgg ttcttgtgtg ttgcccctca agactctctc cctacaaata 240
acttccatat g 251

```

<210> 94

<211> 738

<212> DNA

<213> Homo sapiens

<400> 94

```

cccttttttt tttttttccc acttctcagt ttatttctgg gactaaattt gggtcagagc 60
tgcagagaag ggatgggccc tgagcttggg gatgaaaagt cccagggag attgagacgc 120
aaccctccgcc ctggacagtt ttggaaattg ttcccagggt tcaactagag agacacggtc 180
agcccaattgt gggggaagca gacctgagt ccaggagaca tggggtcagg ggctggagag 240
atgaacatct tcaacatctc tgggaaggaa tgagggtctg aaaggagtgt cagggtctgc 300
cctgcagcag gtggggatgc cggtgtgctg agtcctggga tgactcagga gttggcctgg 360
atggttttct ggtccactt ggtgaacttg cagaggtctg tctagacacc cggtctgttg 420
ggcggggcac aagggtaatc tcccaggac acgagtccct gcaggagacc attgcagacc 480
acaggcccc cagaatcacc ctggcaggag tctctacctg ctttgtcacc ggcgcgaac 540
atggtgtcat ctatctgtct cgggtaagca tctctgcacc ttttctgact tagcacgtg 600
atattcaagc actggaggac cttagggaag tgcacttggg ggctcttggg tgtccccacg 660
ccagacacca agcactttgt cccagcagag ggacaatgag aggagacgtt gatgggtctg 720
acatctttag tgggacga 738

```

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, C12Q 1/68, G01N 33/574, C07K 16/30, A61K 48/00, 31/71, C12N 5/08		A3	(11) International Publication Number: WO 99/33869															
			(43) International Publication Date: 8 July 1999 (08.07.99)															
(21) International Application Number: PCT/US98/27416		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).																
(22) International Filing Date: 22 December 1998 (22.12.98)																		
(30) Priority Data: 08/998,253 24 December 1997 (24.12.97) US 08/998,255 24 December 1997 (24.12.97) US 09/118,627 17 July 1998 (17.07.98) US 09/118,554 17 July 1998 (17.07.98) US																		
(71) Applicant: CORIXA CORPORATION [US/US]: Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).		Published With international search report.																
(72) Inventors: REED, Steven, G.; 2843 - 122nd Place Northeast, Bellevue, WA 98005 (US). XU, Jiangchun; 15805 South-east 43rd Place, Bellevue, WA 98006 (US).		(88) Date of publication of the international search report: 23 December 1999 (23.12.99)																
(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).																		
(54) Title: COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE																		
(57) Abstract																		
<p>Compounds and methods for the treatment and diagnosis of breast cancer are provided. The inventive compounds include polypeptides containing at least a portion of a breast tumor protein. Vaccines and pharmaceutical compositions for immunotherapy of breast cancer comprising such polypeptides, or polynucleotide molecules encoding such polypeptides, are also provided, together with polynucleotide molecules for preparing the inventive polypeptides.</p>																		
<p style="text-align: center;">clone3-6-8</p> <table border="1"> <caption>Data points for clone3-6-8 graph</caption> <thead> <tr> <th>E:T</th> <th>D55LCL/B511s (% specific analysis)</th> <th>D55LCL/HLA-A3 (% specific analysis)</th> </tr> </thead> <tbody> <tr> <td>0.5</td> <td>13</td> <td>0</td> </tr> <tr> <td>1</td> <td>22</td> <td>0</td> </tr> <tr> <td>2</td> <td>27</td> <td>0</td> </tr> <tr> <td>4</td> <td>21</td> <td>0</td> </tr> </tbody> </table>				E:T	D55LCL/B511s (% specific analysis)	D55LCL/HLA-A3 (% specific analysis)	0.5	13	0	1	22	0	2	27	0	4	21	0
E:T	D55LCL/B511s (% specific analysis)	D55LCL/HLA-A3 (% specific analysis)																
0.5	13	0																
1	22	0																
2	27	0																
4	21	0																

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroun			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SU	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

 International Application No.
 PCT/US 98/27416

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/12 C07K14/47 C12Q1/68 G01N33/574 C07K16/30 A61K48/00 A61K31/71 C12N5/08		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 25426 A (CORIXA CORP) 17 July 1997 (1997-07-17) abstract; examples 1-5 page 1, line 1 -page 7, line 11	1-11, 17, 18, 25-34, 36-52
X	WO 97 34921 A (SLOAN KETTERING INST CANCER (US)) 25 September 1997 (1997-09-25) abstract page 7-14 page 22-25 page 31-33	1-3, 7-11, 17, 18
-/-		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (see specification) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
28 June 1999		20.10.99
Name and mailing address of the ISA European Patent Office, P.B. 5018 Patentkan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2000, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Mateo Rosell, A.M.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

 Internatic Application No
 PCT/US 98/27416

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 19783 A (CYTEL CORP (US)) 27 July 1995 (1995-07-27) abstract page 19-28 page 34-35 ---	1-3,17, 18, 25-34, 36-52
X	WO 94 23728 A (UNIV NORTHWESTERN) 27 October 1994 (1994-10-27) page 14, line 25 -page 44, line 2; claim 1; examples 2,3 ---	1-11,17, 18, 25-34, 36-52
X	WO 97 02280 A (HUMAN GENOME SCIENCES INC) 23 January 1997 (1997-01-23) page 1, paragraph 1 page 13, paragraph 2 -page 24, last paragraph ---	1-6, 25-34, 36-46
X	SCHLON J ET AL: "STRATEGIES FOR THE DEVELOPMENT OF RECOMBINANT VACCINES FOR THE IMMUNOTHERAPY OF BREAST CANCER" BREAST CANCER RESEARCH AND TREATMENT, vol. 38, no. 1, 1 January 1996 (1996-01-01), pages 27-39, XP000578043 the whole document ---	1-3,17, 18,47-52
X	PENG LIANG ET AL: "DIFFERENTIAL DISPLAY AND CLONING OF MESSENGER RNAS FROM HUMAN BREAST CANCER VERSUS MAMMARY EPITHELIAL CELLS" CANCER RESEARCH, vol. 52, 15 December 1992 (1992-12-15), pages 6966-6968, XP002032010 the whole document ---	1-3
X	WO 95 11986 A (UNIV COLUMBIA) 4 May 1995 (1995-05-04) page 5, line 1-15; claims 16,21 page 24, line 5-35 page 99-114 ---	1-3
A	K. PORTER-JORDAN AND M.E. LIPPMAN: "Overview of the biological markers of breast cancer" BREAST CANCER, vol. 8, no. 1, 1994, pages 73-100, XP002107232 cited in the application the whole document ---	1
	-/--	

INTERNATIONAL SEARCH REPORT

Internat. Application No.
PCT/US 98/27416

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WATSON M A ET AL: "ISOLATION OF DIFFERENTIALLY EXPRESSED SEQUENCE TAGS FROM HUMAN BREAST CANCER" CANCER RESEARCH, vol. 54, no. 17, 1 September 1994 (1994-09-01), pages 4598-4602, XP000576043 cited in the application the whole document</p>	1
A	<p>DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES, - 3 July 1997 (1997-07-03) XP002106866 HINXTON, GB AC = AA490863. NCI CGAP GCB1 Homo sapiens cDNA clone IMAGE:824150-3'. EST. abstract</p>	1
A	<p>WO 94 21287 A (US HEALTH) 29 September 1994 (1994-09-29) page 1, line 1-6 page 4, line 1 - page 7, line 2; example 1</p>	1-3, 25, 47-52
A	<p>C. YEE ET AL.: "Isolation of tyrosinase-specific CD8+ and CD4+ T cell clones from the peripheral blood of melanoma patients following in vitro stimulation with recombinant vaccinia virus" THE JOURNAL OF IMMUNOLOGY, vol. 157, 1996, pages 4079-4086, XP002106867 cited in the application the whole document</p>	1-3, 17, 18, 47-52

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/27416**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-11, 17, 18, 25-34 and 36-52 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-11,17,18,25-34 and 36-52 all partially

An isolated nucleic acid comprising the nucleotide sequence of SEQ.ID.N.3; encoding an immunogenic portion of a breast protein or a variant of said protein, a vector and a host comprising the nucleic acid, its use for the production of the polypeptide or antibodies reactive with the polypeptide, its use for the manufacture of a vaccine, its use for the detection of breast proteins or nucleic acids.

INTERNATIONAL SEARCH REPORT

int. .ation on patent family members

Internatir Application No

PCT/US 98/27416

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9725426 A	17-07-1997	AU 1697497 A CA 2242340 A CN 1211279 A EP 0874902 A NO 983183 A	01-08-1997 17-07-1997 17-03-1999 04-11-1998 10-09-1998
WO 9734921 A	25-09-1997	AU 2536997 A CA 2249395 A EP 0923605 A	10-10-1997 25-09-1997 23-06-1999
WO 9519783 A	27-07-1995	US 5662907 A AU 1834095 A CA 2181920 A EP 0749315 A SG 49743 A	02-09-1997 08-08-1995 27-07-1995 27-12-1996 15-06-1998
WO 9423728 A	27-10-1994	US 5763164 A AU 6631494 A CA 2160713 A CN 1125399 A EP 0693931 A JP 8509216 T	09-06-1998 08-11-1994 27-10-1994 26-06-1996 31-01-1996 01-10-1998
WO 9702280 A	23-01-1997	CA 2225824 A AU 3092995 A EP 0851869 A JP 11509093 T	23-01-1997 05-02-1997 08-07-1998 17-08-1999
WO 9511986 A	04-05-1995	US 5643761 A AU 695540 B AU 8088394 A EP 0730659 A JP 9507021 T	01-07-1997 13-08-1998 22-05-1995 11-09-1996 15-07-1997
WO 9421287 A	29-09-1994	AU 4002497 A AU 6363094 A CA 2158281 A EP 0692973 A	08-01-1998 11-10-1994 29-09-1994 24-01-1996